

SOLVABILITY IN GROUPS OF PIECEWISE-LINEAR
HOMEOMORPHISMS OF THE UNIT INTERVAL

BY
COLLIN BLEAK

COPY OF DISSERTATION
July, 2005

copyright by

Collin Bleak

2005

Abstract

We investigate subgroups of the group $PL_o(I)$ of piecewise-linear, orientation preserving homeomorphisms of the unit interval with finitely many breaks in slope, and also subgroups of Thompson's group F . We find geometric criteria determining the derived length of any such group, and use this criteria to classify the solvable and non-solvable subgroups of $PL_o(I)$ and of F .

Let H be a subgroup of $PL_o(I)$ or F . We find that H is solvable if and only if H is isomorphic to a group in a well described class $\mathcal{R}$ of groups. We also find that H is non-solvable if and only if we can embed a copy of a specific non-solvable group W into H .

We strengthen the non-solvability classification by finding weak geometric criteria under which we can embed other groups (all containing W) into non-solvable subgroups of $PL_o(I)$ or F .

Acknowledgments

The author would like to acknowledge the patient support of the Mathematics Department at Binghamton University, and the support of his family and friends.

The author further wishes to thank Matthew Brin and Patricia Bleak for their support and advice over the years.

Contents

1	Introduction	1
1.1	An example free statement	1
1.2	Key examples	2
1.3	Formal statements of algebraic results	4
1.4	Geometry	5
1.4.1	Realizing the W_i	5
1.4.2	Some geometric ideas and results	6
2	Classification of solvable subgroups in $PL_o(I)$	8
2.1	Wreath products	8
2.2	The class $\mathcal{R}$	9
2.3	Realizing $\mathcal{R}$ in $PL_o(I)$ and F	13
2.4	$PL_o(I)$	13
2.4.1	Conjugation and orbitals	15
2.4.2	Signed orbitals	16
2.5	Thompson's group F and balanced subgroups of $PL_o(I)$	18
2.5.1	A useful homomorphism	19
2.5.2	Controllers	22
2.6	Towers	25
2.7	Derived groups and towers	31
2.8	Geometric classification of solvable groups in $PL_o(I)$	31
3	Algebraic classification of non-solvable subgroups of $PL_o(I)$	38
3.1	Statement of non-solvability results	39
3.2	Improving exemplary towers	40
3.3	Finding infinite wreath products in groups with infinite towers	50
3.4	W and other groups	60
3.5	W in arbitrary non-solvable subgroups of $PL_o(I)$	63

1 Introduction

We find descriptions of solvable and non-solvable subgroups of $PL_o(I)$, the group of orientation preserving piecewise linear homeomorphisms of the unit interval with finitely many breaks in slope. One description we find of the solvable subgroups of $PL_o(I)$ is as the set of isomorphism classes of the subgroups of an easily described, countable set $\mathcal{M} = \{G_0, G_1, G_2, \dots\}$ of countable, solvable subgroups of $PL_o(I)$. We also find one non-solvable group W so that a subgroup H of $PL_o(I)$ is non-solvable if and only if H contains an isomorphic copy of W . We will say more about these descriptions, and others, later on in the introduction.

Much of our analysis of the solvable and non-solvable subgroups of $PL_o(I)$ does not rely on the piecewise linear nature of the homeomorphisms of the unit interval; we tend to use point “dynamics” under homeomorphisms of the interval. It would be interesting to discover how close the classifications below come to classifying the solvable and non-solvable subgroups of $Homeo_+(I)$, the group of orientation preserving homeomorphisms of the unit interval.

Unknown to the author at the time of this work, Andrés Navas ([5], [6]) has also been using dynamics to analyze the solvable subgroups of the group $Diff_+^2(\mathbf{R})$. In [6], he uses his results to show that a finitely generated solvable subgroup of $PL_o(I)$ with connected support is isomorphic to a semi-direct product of a group H with the integers $\mathbf{Z}$, where H is a group in a special class of groups (we also discover this class of groups). This result is contained in our investigations below, but his techniques are different from our own. Perhaps a combination of Navas’ techniques with those below may be sufficient to classify the solvable and non-solvable subgroups of $Homeo_+(I)$.

Our investigations also have an impact on the theory of Thompson’s group F . Both the class $\mathcal{M}$ of groups that we realize in $PL_o(I)$, and the group W , can be realized in the standard realization of F in $PL_o(I)$, so that we also have descriptions of the solvable and non-solvable subgroups of F .

1.1 An example free statement

Most of our understanding of the subgroups of $PL_o(I)$ is derived through analysis of important examples of subgroups of $PL_o(I)$. Nonetheless, we give here an example free description of the solvable subgroups of $PL_o(I)$.

Define $\mathcal{R}$ to be the smallest non-empty class of groups which is closed under the following three operations:

1. Restricted wreath product with $\mathbf{Z}$ (ie. $H \mapsto H \wr \mathbf{Z}$).
2. Bounded direct sum (defined below).
3. Taking subgroups.

Here the “bounded direct sum” is a countable direct sum of groups in $\mathcal{R}$ for which there is a uniform bound on the lengths of their derived series.

With this definition in place, we can give another algebraic version of our result on the solvable groups in $PL_o(I)$:

Theorem 1.1 *H is isomorphic to a solvable subgroup of $PL_o(I)$ if and only if H is isomorphic to a group in the class $\mathcal{R}$ of groups.*

1.2 Key examples

In this section, we will mention some key examples. All of these examples can be realized in $PL_o(I)$, and even in Thompson’s group F . Our examples rely on an understanding of the wreath product, both as a standard restricted wreath product as discussed by P. M. Neumann in [7], and as a permutation wreath product which is discussed in detail in section 2.1. In situations such as

$$(\dots((G_1 \wr G_2) \wr G_3) \wr \dots \wr G_n)$$

where parentheses “accumulate on the left” and each G_i acts on itself by right multiplication, the resulting groups are the same if $\wr$ represents the standard restricted wreath product or the permutation wreath product. Until we need to distinguish the two types of wreath product we will not do so.

Before progressing into this discussion, let us fix $\mathbf{N}$ as representing the positive integers for the remainder of our investigations.

First, let us build the class $\mathcal{M}$ of groups. Define

$$G_0 = \{1\},$$

the trivial group, and for each $n \in \mathbf{N}$, inductively define

$$G_n = \bigoplus_{i \in \mathbf{Z}} (G_{n-1} \wr \mathbf{Z}).$$

Note, for example, that $G_1 \cong \bigoplus_{i \in \mathbf{Z}} \mathbf{Z}$. Now define

$$\mathcal{M} = \{G_i \mid i \in \mathbf{Z}, i \geq 0\}.$$

The class $\mathcal{M}$ of groups has nice properties, the first of which is obvious, and the latter two of which we will prove later:

1. Given non-negative $i \in \mathbf{Z}$, $G_i \cong \bigoplus_{j \in \mathbf{Z}} G_i$ (note: the subscript is not the sum index).

2. Given non-negative $n \in \mathbf{Z}$, G_n has derived length n .
3. If H is a subgroup of G_k for some non-negative $k \in \mathbf{Z}$, and H has derived length n , then H is isomorphic to a subgroup of G_n .

Our main interest in these groups is that they play a key role in understanding the class $\mathcal{R}$.

We will see that taking a restricted wreath product with $\mathbf{Z}$ is something that is easy to realize in $PL_o(I)$, which is why this activity plays a key role in the definition of the class $\mathcal{M}$. In fact, it is so natural that another collection of groups becomes relevant. Define $W_0 = 1$, the trivial group, and for all $i \in \mathbf{N}$, define $W_i = W_{i-1} \wr \mathbf{Z}$, so that

$$W_i = (\dots (((\mathbf{Z} \wr \mathbf{Z}) \wr \mathbf{Z}) \wr \mathbf{Z}) \dots) \wr \mathbf{Z},$$

where there are i appearances of $\mathbf{Z}$ on the right. Our main result towards understanding $\mathcal{R}$ can be rephrased in terms of the W_i instead of the G_i . However, the W_i lack the corresponding first and third properties of the G_i stated above, and these are serious deficiencies in the class, from a computational point of view.

Nonetheless, the W_i are useful since an isomorphic copy of the group

$$W = \bigoplus_{i \in \mathbf{N}} W_i$$

occurs as a subgroup of any non-solvable subgroup of $PL_o(I)$.

We now describe some often recurring groups, all of which contain an isomorphic copy of W as a subgroup. The first group is

$$(\mathbf{Z})^\infty = \dots (((\mathbf{Z} \wr \mathbf{Z}) \wr \mathbf{Z}) \wr \mathbf{Z}) \wr \dots$$

and the second group is the permutation wreath product:

$$(\wr \mathbf{Z})^\infty = \dots \wr (\mathbf{Z} \wr (\mathbf{Z} \wr (\mathbf{Z} \wr \mathbf{Z}))) \wr \dots$$

Sapir had raised the question of whether every non-solvable subgroup of F contained an isomorphic copy of $(\mathbf{Z})^\infty$. In [2], Brin shows that both $(\mathbf{Z})^\infty$ and $(\wr \mathbf{Z})^\infty$ occur as non-solvable subgroups of Thompson's group F , but that neither $(\mathbf{Z})^\infty$ nor $(\wr \mathbf{Z})^\infty$ contain the other as a subgroup. Brin also shows that a third group, the permutation wreath product

$$(\wr \mathbf{Z})^\infty = (\wr \mathbf{Z})^\infty \wr (\mathbf{Z})^\infty$$

is also realized in F . The group $(\wr \mathbf{Z})^\infty$ contains both an isomorphic copy of $(\mathbf{Z})^\infty$ and an isomorphic copy of $(\wr \mathbf{Z})^\infty$. A natural follow-up question to Sapir's initial question is whether one of $(\mathbf{Z})^\infty$ or $(\wr \mathbf{Z})^\infty$ is always to be found as a subgroup of any non-solvable subgroup of F . These investigations answer this question. See the next section.

1.3 Formal statements of algebraic results

There will be a later section giving geometric results which are used to obtain the results we give here. In this section, we will state all of our main results, and make a few comments.

Our two chief results are Theorem 1.1 above, and the following:

Theorem 1.2 *Suppose H is a subgroup of $PL_o(I)$. H is non-solvable if and only if H contains a subgroup isomorphic to W .*

Theorem 1.1 is further explained by the following result:

Theorem 1.3 *$H \in \mathcal{R}$ if and only if H is isomorphic to a subgroup of a group in $\mathcal{M}$.*

This second description of the class $\mathcal{R}$ of groups will greatly assist us in doing calculations relating to the solvable subgroups of $PL_o(I)$ from a purely algebraic perspective. The proofs of both Theorem 1.1 and Theorem 1.3 will depend on the following Lemma.

Lemma 1.4 *If G is a solvable subgroup of $PL_o(I)$ with derived length n , then G is isomorphic to a subgroup of G_n .*

It is immediate from construction that the groups in $\mathcal{M}$ are all countable, so the last lemma also has the following corollary.

Corollary 1.5 *If H is a solvable subgroup of $PL_o(I)$, then H is countable.*

Since the groups in $\mathcal{M}$ can all be realized in Thompson's group F , we further have:

Corollary 1.6 *A group $H \in \mathcal{R}$ if and only if H is isomorphic with a solvable subgroup of F .*

It is easily seen that W admits no finite index solvable subgroup, so we have another corollary based on Theorem 1.2.

Corollary 1.7 *Virtually solvable subgroups of $PL_o(I)$ are solvable.*

Theorem 1.2 does not tell the whole story of the non-solvable subgroups of $PL_o(I)$. Later we give weak geometric conditions, each of which implies that a subgroup of $PL_o(I)$ contains a subgroup isomorphic to $(\mathbb{Z})^\infty$.

We also show the following:

Theorem 1.8 *If H is a finitely generated non-solvable subgroup of $PL_o(I)$, then H contains an isomorphic copy of $(\mathbb{Z})^\infty$ or $(\mathbb{Z})^\infty$.*

It is a question whether every finitely generated non-solvable subgroup of $PL_o(I)$ contains $(\mathbb{Z})^\infty$. (Note, W is not finitely generated and non-solvable, and no copy of $(\mathbb{Z})^\infty$ embeds in it.)

1.4 Geometry


In this section, we will realize the W_i in $F \leq PL_o(I)$, and use these realizations to motivate some geometric definitions which will enable us to state our main geometric result upon which both Theorem 1.1 and Theorem 1.2 depend. Here, we are thinking of F as the realization of Thompson's group in $PL_o(I)$ which consists of all the elements of $PL_o(I)$ which have all slopes powers of two, and which have all breakpoints occurring at the dyadic rationals $\mathbf{Z}[\frac{1}{2}]$. See Cannon, Floyd, and Parry [4] for an introduction to the remarkable group F .

1.4.1 Realizing the W_i

Consider the two elements $\alpha_1, \alpha_2 \in PL_o(I)$ defined below:

$$x\alpha_1 = \begin{cases} 2x & 0 \leq x < \frac{1}{4}, \\ x + \frac{1}{4} & \frac{1}{4} \leq x < \frac{1}{2}, \\ \frac{1}{2}x + \frac{1}{2} & \frac{1}{2} \leq x \leq 1, \end{cases} \quad x\alpha_2 = \begin{cases} x & 0 \leq x < \frac{1}{4}, \\ 2x - \frac{1}{4} & \frac{1}{4} \leq x < \frac{5}{16}, \\ x + \frac{1}{16} & \frac{5}{16} \leq x \leq \frac{3}{8}, \\ \frac{1}{2}x + \frac{1}{4} & \frac{3}{8} \leq x < \frac{1}{2}, \\ x & \frac{1}{2} \leq x \leq 1. \end{cases}$$

Here are the graphs (superimposed) of these functions:



Either element alone generates a group isomorphic to $\mathbf{Z} \cong W_1$ in $PL_o(I)$, but the “action” of α_2 occurs in a single fundamental domain of α_1 ; that is, $\frac{1}{4}\alpha_1 = \frac{1}{2}$, but α_2 is the identity

off of the interval $[\frac{1}{4}, \frac{1}{2}]$. In particular, $\alpha_2^{\alpha_1} = \alpha_1^{-1}\alpha_2\alpha_1$ has support $(\frac{1}{2}, \frac{3}{4})$, which is disjoint from the support of α_2 . (In this discussion, following the notation in Brin's papers [1] and [2], elements of $PL_o(I)$ act on the right on I , and the support of any particular element of $PL_o(I)$ is the open set of points in I that are moved by that element.) In particular, any two distinct conjugates of α_2 by powers of α_1 commute with each other, since their supports will be disjoint in the interval I . Now, if we consider any element $h \in \langle \alpha_1, \alpha_2 \rangle$, it is a standard algebraic fact that we can write h as a product of the form a power of α_1 followed by a product of conjugates of α_2 (and α_2^{-1}) by various powers of α_1 . In particular, the element α_1 generates a group isomorphic to $\mathbf{Z}$ which acts on the normal subgroup of $\langle \alpha_1, \alpha_2 \rangle$ consisting of the direct sum of $\mathbf{Z}$'s generated by the conjugates of α_2 by different powers of α_1 . The following chain of isomorphisms should now make sense:

$$\langle \alpha_1, \alpha_2 \rangle \cong \left(\bigoplus_{i \in \mathbf{Z}} \mathbf{Z} \right) \rtimes \mathbf{Z} \cong \mathbf{Z} \wr \mathbf{Z} \cong W_2.$$

Note before we move on that α_1 and α_2 are both elements of Thompson's group F .

All of the W_i can be realized in Thompson's group F in an entirely similar way, using one-bump generators; take α_i to be the conjugate $\alpha_i = \alpha_{i-1}^s = s^{-1}\alpha_{i-1}s$, where s is the "shrinking function" $s(x) = \frac{1}{4}x + \frac{1}{4}$, and the conjugation takes place in $PL_o(\mathbf{R})$, where we replace α_1 by the function in $PL_o(\mathbf{R})$ which behaves as the identity outside of the unit interval $[0, 1]$ for this inductive definition. Given $i \in \mathbf{N}$, the support of α_i is contained in a single fundamental domain of α_{i-1} , so that given $j \in \mathbf{N}$, we have $W_j \cong \langle \alpha_1, \alpha_2, \dots, \alpha_j \rangle$. Since conjugating any element of F by s will still produce an element of F , we see that the W_i 's can all be realized in F .

1.4.2 Some geometric ideas and results

Much of the language of this section is motivated by thinking of subgroups of $PL_o(I)$ as permutation groups acting on the set $I = [0, 1]$. We will become progressively less formal in this section, as we are only trying to indicate the nature of our geometric results. Formal definitions will be given in later sections.

If H is a subgroup of $PL_o(I)$, then its support naturally falls into a collection of disjoint, open intervals. Each such we will call an *orbital* of the group H . Given an element $\gamma \in H$, the group $\langle \gamma \rangle$ has its own orbitals, which are the connected components of the support of γ . Given such an interval $A = (a, b) \subset [0, 1]$, we call A an *orbital* of γ . Since H now must have infinitely many elements that also have A as an orbital, we typically use *signed orbitals*, which are pairs of the form (A, γ) to indicate not only which element orbital we are considering, but also the specific element which is "owning" that orbital in our discussion. In the example groups W_i , each of the α_k had an orbital whose closure was fully contained in the orbital of α_{k-1} , whenever $k > 1$. Such a "stack" of k signed, nested orbitals forms what we call a "tower". Observe that many subgroups of $PL_o(I)$ have pairs of elements with orbitals that overlap each other on one end, and are therefore not arranged as a tower. In this case, we can use the respective elements to move points in I across one orbital into one end of an overlapping orbital, and then use the second element to move the points across the new orbital out of the original orbital. This horizontal technique, and arrangement of orbitals,

motivates our definition of a “transition chain”, which consists of a “chain” of neighbor-overlapping signed orbitals. Typically, the elements of a tower generate easier groups to understand, while the set of elements of a transition chain generate highly complex groups.

We are now in a position to approach our main geometric result. We will say that a group has depth $n \in \mathbf{N}$ if and only if we can find towers of height n , but no towers of height $n + 1$, in the group. Our main geometric result is as follows:

Theorem 1.9 *Suppose G is a subgroup of $PL_o(I)$ and $n \in \mathbf{Z}$ with $n \geq 0$. G is solvable with derived length n if and only if G has depth n .*

Theorem 1.9 is used in both the classification of the solvable subgroups of $PL_o(I)$ and the classification of the non-solvable subgroups of $PL_o(I)$.

As hinted at above, the techniques we use can naturally be broken down into two types: vertical techniques that correspond to working with towers, and horizontal techniques that correspond to working with transition chains. Vertical techniques lead to Theorem 1.9. The horizontal techniques extract information from the transition chains, and then work to get rid of them.

2 Classification of solvable subgroups in $PL_o(I)$

In this section we will pursue the algebraic classification of the solvable subgroups in $PL_o(I)$. One direction of the classification is mostly algebraic, and requires less knowledge of the terminology of $PL_o(I)$. We will engage in that direction first.

2.1 Wreath products

In this paper the restricted wreath product is as given by Neumann in [7]. The restricted wreath product is distinguished from the unrestricted wreath product in that the restricted wreath product $A \wr B$ is formed as $C \rtimes B$ where C is a sum of copies of A as opposed to a product of copies of A . One can also think of a restricted wreath product of two groups as the group resulting from a permutation wreath product (definition below) where the sets that the two groups in the product act on are their own underlying sets using right multiplication.

In [8], Robinson describes the permutation wreath product, and proves some of the basic facts about this product. We will generally follow his presentation, with only minor modifications.

Let H and K be permutation groups acting on sets X and Y respectively. We will describe how to construct a new permutation group $H \wr K$, called the permutation wreath product of H and K . Our new group $H \wr K$ will act on the set $Z = X \times Y$.

If $\gamma \in H$, $\kappa \in K$, and $y \in Y$, define the permutations γ_y and κ^* of Z by the rules:

$$\gamma_y : \begin{cases} (x, y) \mapsto (x\gamma, y), \\ (x, y') \mapsto (x, y') \text{ if } y' \neq y, \end{cases}$$

$$\kappa^* : (x, y) \mapsto (x, y\kappa).$$

We note that $(\gamma^{-1})_y = (\gamma_y)^{-1}$ and $(\kappa^{-1})^* = (\kappa^*)^{-1}$, so that γ_y and κ^* are invertible and therefore really are elements in $Sym(Z)$ (the group of permutations of the set Z). One can check that with y fixed, the map $\phi_y : H \rightarrow Sym(Z)$ defined by the rule $\gamma \mapsto \gamma_y$ is a monic homomorphism, and regardless, that $\phi^* : K \rightarrow Sym(Z)$ defined by the rule $\kappa \mapsto \kappa^*$ is also a monic homomorphism. (The monic property follows from the definition of permutation groups as subgroups of the symmetric group on the set being acted upon, so that all elements except the identity move something.) If we denote the images of ϕ_y and ϕ^* by H_y and K^* respectively, then *the wreath product of H and K is the subgroup of $Sym(Z)$ generated by all of the H_y for $y \in Y$ and the group K^** , ie. $H \wr K = \langle H_y, K^* \mid y \in Y \rangle$.

Because the γ_y have disjoint support in Z , the H_y 's generate a subgroup B of $H \wr K$ with $B \cong \bigoplus_{y \in Y} H_y$. The group B is normal in $H \wr K$, since $(\kappa^*)^{-1} \gamma_y \kappa^*$ acts by mapping $(x, y\kappa)$ to $(x\gamma, y\kappa)$, and by fixing (x', y') if $y' \neq y\kappa$, ie., $(\kappa^*)^{-1} \gamma_y \kappa^* = \gamma_{y\kappa} \in B$. In particular, conjugating B by an element κ^* of K^* simply permutes the direct factors of B the same way that κ permutes the elements of Y . Now since non-trivial elements of B always move some element of Z , and can only effect the first coordinate of such elements of Z , while non-trivial elements of K^* always move some element of Z , and can only effect the second coordinates of such elements, we see that $B \cap K^* = 1$ in $H \wr K$, so that $H \wr K \cong B \rtimes K^*$. We typically call B the “Base group” of $H \wr K$, and K^* (or K for simplicity) the “Top group” of $H \wr K$. We will often use the coordinate representation (or implied variants of it) of $H \wr K$ as the set of elements of the form (b, k) where $b \in B$ and $k \in K$ in our arguments. We will often denote a $b \in \bigoplus_{y \in Y} H_y$ by $(b)_{y \in Y}$.

As mentioned in the introduction, we need a little care to specify when we are using the restricted wreath product or when we are using the permutation wreath product. Here is an example: If we temporarily let $\wr_p$ represent the permutation wreath product, and $\wr$ represent the restricted wreath product, and assume that $\mathbf{Z}$ is a permutation group acting on its base set by right multiplication in the permutation wreath products below, then observe the difference in the following two groups:

$$\begin{aligned} A &= \mathbf{Z} \wr (\mathbf{Z} \wr \mathbf{Z}) \cong (\bigoplus_{a \in \mathbf{Z} \wr \mathbf{Z}} \mathbf{Z}) \rtimes \mathbf{Z} \wr \mathbf{Z}, & \text{and} \\ B &= \mathbf{Z} \wr_p (\mathbf{Z} \wr_p \mathbf{Z}) \cong (\bigoplus_{a \in \mathbf{Z} \times \mathbf{Z}} \mathbf{Z}) \rtimes \mathbf{Z} \wr_p \mathbf{Z}. \end{aligned}$$

The top groups are isomorphic, and so are the base groups, but overall the groups are very different in the sense of how the summands of the base group are indexed and moved by the action of the top group.

Note that differences arise in the two wreath products (restricted and permutation) since in the permutation wreath product, the set associated with the group may not actually be the underlying set of the group. When we take wreath products on the right with $\mathbf{Z}$, this will never be an issue, so we will not focus on this issue unless it needs explicit care (such as in the construction of $(\wr \mathbf{Z})^\infty$).

Note that in [8], Robinson shows that the permutation wreath product of any three permutation groups is associative. From the last example, we see that this is not the case for restricted wreath products.

One should consider the realization of W_2 in the introduction to understand what sets play the roles of X , Y , and Z , and what are the incarnations of the groups B and K in that context. Brin’s paper [2] goes in depth into understanding this realization.

2.2 The class $\mathcal{R}$

Recall that $\mathcal{R}$ represents the smallest non-empty class of groups which is closed under the following three operations.

1. Restricted wreath product with $\mathbf{Z}$.
2. Bounded direct sum.

3. Taking subgroups.

Here bounded direct sum means a direct sum of groups in $\mathcal{R}$, all of whose derived lengths are less than some bounding integer M , and where the index set of the direct sum is countable.

We will investigate this class and come to understand it from a second perspective, and then by using Theorems 1.3 and 1.9, we will prove Theorem 1.1.

We begin to understand $\mathcal{R}$ via a close study of the class $\mathcal{M} = \{G_i \mid i \geq 0, i \in \mathbf{Z}\}$ defined in the introduction. Let us gather some facts about the groups G_k .

Lemma 2.1 *1. If F_0, F_1, H_0 , and H_1 are groups, where $F_0 \leq F_1$ and $H_0 \leq H_1$, then $F_0 \wr H_0 \leq F_1 \wr H_1$.*

2. For any non-negative m and $n \in \mathbf{Z}$, with $m < n$, G_m embeds as a normal subgroup of G_n .

3. For any group G with derived length n , the groups $G \wr \mathbf{Z}$ and $\bigoplus_{i \in \mathbf{Z}} (G \wr \mathbf{Z})$ have derived length $n + 1$.

4. For any non-negative $n \in \mathbf{Z}$, G_n has derived length n .

pf: The first point is immediate by examining the following chain of subgroup inclusions, where the inclusions are based on the underlying sets.

$$\begin{aligned} F_0 \wr H_0 &= \{((f_0)_{h \in H_0}, h_0) \mid (f_0)_{h \in H_0} \in \bigoplus_{a \in H_0} F_0, h_0 \in H_0\} \leq \\ F_1 \wr H_0 &= \{((f_1)_{h \in H_0}, h_0) \mid (f_1)_{h \in H_0} \in \bigoplus_{a \in H_0} F_1, h_0 \in H_0\} \leq \\ F_1 \wr H_1 &= \{((f_1)_{h \in H_1}, h_1) \mid (f_1)_{h \in H_1} \in \bigoplus_{a \in H_1} F_1, h_1 \in H_1\} \end{aligned}$$

To see the second point, we will demonstrate an embedding of G_{n-1} into G_n , and thus inductively define an embedding of G_m into G_n , for any non-negative integers $m < n$. Note that there are many copies of G_{n-1} in G_n , but we are particularly interested in the one given in the next paragraph, which is the copy that we use to inductively define our particular copy of G_m in G_n . The normality of this embedded copy of G_m in G_n follows easily from the theory of group actions, which can be checked in section 2.3 below where we realize G_n in $PL_o(I)$. (Let X be the set of left hand endpoints of the components of the support of G_m in I . X is acted upon by G_n , and the kernel of this action is G_m .) A second proof is by noting that the embedded copy of G_{n-1} in G_n that we demonstrate in the paragraph below is characteristic in G_n , which proof can be carried out with the help of the geometric tools established in the proof of Theorem 1.9. We will not use the normality of our embedded copy of G_m in G_n later.

Now let us describe our particular embedded copy of G_{n-1} in G_n . First, identify the base group of $G_{n-1} \wr \mathbf{Z}$ with G_{n-1} using the fact that $\bigoplus_{i \in \mathbf{Z}} G_{n-1} \cong G_{n-1}$, now since the direct sum of the base groups of the $G_{n-1} \wr \mathbf{Z}$ summands in the definition of G_n is also a subgroup of G_n , we see that $\bigoplus_{i \in \mathbf{Z}} G_{n-1}$ is a subgroup of G_n . But now again, this last direct sum is isomorphic with G_{n-1} , so that G_{n-1} (as embedded here as the direct sum of the base groups of the $G_{n-1} \wr \mathbf{Z}$ summands of G_n) is a subgroup of G_n .

For the third point, it follows from Neumann [7] (Theorem 4.1 and Corollary 4.5) that if $G = A \wr B$, then G' is contained in a sum of copies of A , and also that G' surjects onto A (these facts are under the condition that B is abelian). Both facts are easy exercises in our situation with $B = \mathbf{Z}$. This immediately implies the third point.

The fourth point follows directly from the third.

◇

Now let us examine $\mathcal{R}$ for a short time. We give (modulo work to come later) a characterization of $\mathcal{R}$ that completes one direction of Theorem 1.1. The key result (Lemma 2.4 below) will not be used in the rest of the paper.

Since $\mathcal{R}$ is nonempty and closed under subgroups, the group 1 is a group in $\mathcal{R}$. Let $\mathcal{G}$ represent the class of all groups. Define $\mathcal{P} : \mathcal{P}(\mathcal{G}) \rightarrow \mathcal{P}(\mathcal{G})$ to be the function representing the closure operation that takes a set X of groups and computes the smallest class of groups which contains X and is closed under the operations of restricted wreath product with $\mathbf{Z}$ and bounded direct sum. Since each G_i is obtained by applying a finite sequence of bounded direct sums and wreath products with $\mathbf{Z}$ to the trivial group 1, we see that $\mathcal{M} \subset \{1\}\mathcal{P}$. By the definition of $\mathcal{R}$, it is immediate that $\{1\}\mathcal{P} \subset \mathcal{R}$. In particular, we have:

$$\mathcal{M} \subset \{1\}\mathcal{P} \subset \mathcal{R}$$

Now let us consider three operators

$$\begin{aligned} \mathcal{S} : \mathcal{P}(\mathcal{G}) &\rightarrow \mathcal{P}(\mathcal{G}), \\ \mathcal{W} : \mathcal{P}(\mathcal{G}) &\rightarrow \mathcal{P}(\mathcal{G}), \\ \mathcal{B} : \mathcal{P}(\mathcal{G}) &\rightarrow \mathcal{P}(\mathcal{G}), \end{aligned}$$

that represent taking closure under the operations of taking subgroups, taking restricted wreath products with $\mathbf{Z}$, and building bounded direct sums, respectively.

We now investigate the relationship between the closure operations of the three actions; taking subgroups, computing restricted wreath products with $\mathbf{Z}$, and computing bounded direct sums. If Γ is the set of all finite length words of the form $(\mathcal{WB})^k$ or $(\mathcal{BW})^k$ where $k \in \{0, 1, 2, \dots\}$, and $X \in \mathcal{P}(\mathcal{G})$, then the union $\Upsilon_X = \cup_{\gamma \in \Gamma} X\gamma$ is the smallest closed set containing X that is closed under both operations of taking bounded direct sums and wreath products with $\mathbf{Z}$, so that $\Upsilon_{\{1\}} = \{1\}\mathcal{P}$. In particular, the smallest class of groups which contains the trivial group and which is closed under the operations of taking wreath products with $\mathbf{Z}$ and building bounded direct sums equals $\{1\}\mathcal{P}$.

Lemma 2.2 *Let m be a non-negative integer. If G is a group in $\{1\}\mathcal{P}$ with derived length m , then G is isomorphic to a subgroup of G_m .*

pf: We can prove this by inducting on the derived length of G . If G has derived length 0, then G is the trivial group so $G = G_0$. Let $n \in \mathbf{N}$, and suppose G has derived length n and that for any group H in $\{1\}\mathcal{P}$ which has derived length m where $0 \leq m < n$, we know that $H \leq G_m$. Now, there is a $j \in \mathbf{N}$ so that $G \in \{1\}(\mathcal{BW})^j$, or $G \in \{1\}(\mathcal{WB})^j$. Note that if $G \in \{1\}(\mathcal{WB})^j$ then $G \in \{1\}(\mathcal{BW})^{j+1}$, so we will assume that $G \in \{1\}(\mathcal{BW})^k$ for some minimal non-negative integer k . There are now two cases:

1. $G \in \{1\}(\mathcal{BW})^{k-1}\mathcal{B}$ but G is not in $\{1\}(\mathcal{BW})^{k-1}$.

In this case, The last operation required to build G was a bounded direct sum of groups, all of which groups have derived lengths necessarily less than or equal to n (note here that a finite sequence of bounded direct sums is isomorphic to a bounded direct sum). The summands of this bounded direct sum are all groups in $\{1\}(\mathcal{BW})^{k-1}$. We will argue that each of these groups is actually a subgroup of G_n , and therefore, since a countable or finite direct sum of groups isomorphic to G_n is actually isomorphic to G_n , we will have finished this case.

Let H be a summand of the final bounded direct sum which created G , and assume that the derived length of H is actually n (at least one summand must have this derived length), and that $H \in \{1\}(\mathcal{BW})^{k-1}$. If H is actually in $\{1\}(\mathcal{BW})^{k-2}\mathcal{B}$, then we can replace H inductively by a summand (with derived length n) of the last bounded sum operation used to create H , so that there is a $t \in \mathbf{N}$ so that H is now an element of $\{1\}(\mathcal{BW})^t$, but not an element of $\{1\}(\mathcal{BW})^{t-1}\mathcal{B}$. In particular, H is the result of applying s wreath products with $\mathbf{Z}$ to a group H^* in $\{1\}(\mathcal{BW})^{t-1}\mathcal{B}$ for some $s \in \mathbf{N}$. H^* is therefore a group in $\{1\}\mathcal{P}$ with derived length $n - s$, and therefore H^* is a subgroup of G_{n-s} . But note that for any integer p we have that $G_p \wr \mathbf{Z} \leq G_{p+1}$, so that $H = H^*(\wr \mathbf{Z})^s \leq G_{n-s}(\wr \mathbf{Z})^s \leq G_{n-s+1}(\wr \mathbf{Z})^{s-1} \leq \dots \leq G_n$.

If H is a summand of G with derived length m where $m < n$, Then by our induction hypothesis, $H \leq G_m$, but $G_m \leq G_n$, so $H \leq G_n$.

2. $G \in \{1\}(\mathcal{BW})^k$ but G is not in $\{1\}(\mathcal{BW})^{k-1}\mathcal{B}$.

This case is entirely similar to the last case, except that we already know that G is the result of applying s wreath products with $\mathbf{Z}$ to a group H in $\{1\}(\mathcal{BW})^{k-1}\mathcal{B}$ for some positive integer s . The derived length of H must be $n - s$, and therefore $H \leq G_{n-s}$ so that $G \leq G_n$ as in the penultimate paragraph of the previous case.

◇

We need one last technical lemma before we can complete our exploration of the class $\mathcal{R}$:

Lemma 2.3 *Suppose m and $n \in \mathbf{Z}$, with $0 \leq m \leq n$. If $H \leq G_n$ and H has derived length m , then H is isomorphic to a subgroup of G_m .*

pf: This follows from Corollary 2.36 below, and the fact that we can realize the groups $\mathcal{M}$ in $PL_o(I)$ (see next subsection). ◇

Our arguments after this section do not rely on the classification of $\mathcal{R}$ given in the next lemma.

Finally, we have a nice description of $\mathcal{R}$. Note that the following lemma implies Theorem 1.3.

Lemma 2.4 $\mathcal{R} = \{1\}\mathcal{PS} = \mathcal{MS}$

Pf: We have already shown that $\{1\}\mathcal{P} \subset \mathcal{MS}$, so we know that $\{1\}\mathcal{PS} \subset \mathcal{MS}$, and by definition, $\mathcal{M} \subset \{1\}\mathcal{P}$, so that $\mathcal{MS} \subset \{1\}\mathcal{PS}$. In particular, $\mathcal{MS} = \{1\}\mathcal{PS}$.

We will now show that $\mathcal{MSB} = \mathcal{MS}$ (implying that $\mathcal{MS}$ is already closed under the operation of taking bounded direct sums) and that $\mathcal{MSW} = \mathcal{MS}$ (implying that $\mathcal{MS}$ is already closed under wreath products with $\mathbf{Z}$, so that we can conclude that $\mathcal{MS} = \mathcal{R}$).

To see that $\mathcal{MSB} = \mathcal{MS}$, let $G \in \mathcal{MSB}$. There is an $M \in \mathbf{N}$ so that we can write $G = \bigoplus_{i \in \mathbf{Z}} H_i$, where each H_i has derived length bounded above by M . Now, each $H_i \leq G_M$ (by the Lemma 2.3), so we see that $G \leq \bigoplus_{i \in \mathbf{Z}} G_M$, hence $G \in \mathcal{MS}$.

To see that $\mathcal{MSW} = \mathcal{MS}$, we note that if $G \in \mathcal{MSW}$, then either $G \in \mathcal{MS}$ or we can write $G = ((\dots((H \wr \mathbf{Z}) \wr \mathbf{Z}) \dots) \wr \mathbf{Z})$, where $H \in \mathcal{MS}$, and where there are k wreath products with $\mathbf{Z}$, for some $k \in \mathbf{N}$. In the first case we are done. In the second case $H \leq G_M$ for some non-negative integer M . But now $G = ((\dots((H \wr \mathbf{Z}) \wr \mathbf{Z}) \dots) \wr \mathbf{Z}) \leq ((\dots((G_M \wr \mathbf{Z}) \wr \mathbf{Z}) \dots) \wr \mathbf{Z}) = J \in \{1\} \mathcal{P}$, where J has derived length $M + k$, so that $G \leq G_{M+k}$. Finally we have that $G \in \mathcal{MS}$. $\diamond$

2.3 Realizing $\mathcal{R}$ in $PL_o(I)$ and F

Here we will explain how we can realize the groups in $\mathcal{M}$ inside Thompson's group F , as realized in $PL_o(I)$. This will prove one half of Theorem 1.1. To realize the other direction of Theorem 1.1, we will need to take a lengthy detour through the geometric definitions of $PL_o(I)$.

The elements $\alpha_1, \alpha_2 \in PL_o(I)$ defined in the introduction will play a major role here, as will the shrinking conjugator s .

First, observe that we can realize G_1 fairly simply. Let $\beta_0 = \alpha_2$, and define β_k for each $k \in \mathbf{Z}$ as $\beta_0^{\alpha_1^k}$. G_1 is immediately isomorphic to $\langle \beta_k | k \in \mathbf{Z} \rangle$, and G_1 has been realized in Thompson's group F .

Now we will show that given any group H which has been realized as a subgroup of $PL_o(I)$, we can realize $\bigoplus_{i \in \mathbf{Z}} (H \wr \mathbf{Z})$ as a subgroup of $PL_o(I)$. Firstly, conjugate the elements of H by s twice, to create a new group H_0 isomorphic with H . The supports of all of the elements of H_0 are contained in the set $(\frac{5}{16}, \frac{3}{8})$, which is contained in a single fundamental domain of $\alpha_2 = \beta_0$. At this juncture, the group generated by H_0 and β_0 is isomorphic to $H \wr \mathbf{Z}$, where β_0 is the generator of the top $\mathbf{Z}$ factor. In particular, we can realize $H \wr \mathbf{Z}$ in $PL_o(I)$. But now $\bigoplus_{i \in \mathbf{Z}} (H \wr \mathbf{Z})$ is the base group of $(H \wr \mathbf{Z}) \wr \mathbf{Z}$, which we can also realize by repeating the previous procedure, so we are done.

Observe that the shrinking map conjugates elements of Thompson's group F into Thompson's group F , so that if H is realized as a subgroup of F , then this construction of $\bigoplus_{i \in \mathbf{Z}} (H \wr \mathbf{Z})$ also produces a subgroup of F . In particular, we see inductively that each group G_n can now be realized in Thompson's group F , and therefore that each group of $\mathcal{R}$ can be realized in Thompson's group F .

2.4 $PL_o(I)$

Beginning with this section, and through to the end of section 2.6, we build the geometric tools and analysis necessary to complete the proof of Theorem 1.1.

We will now build some required terminology for working with subgroups of $PL_o(I)$. We will use notation similar to that in Brin's paper [1] on the ubiquitous nature of Thompson's group F in subgroups of $PL_o(I)$.

We note that the set of points moved by an element h of $PL_o(I)$ is open, by the continuity of elements of $PL_o(I)$. But then the support of H , for any subgroup $H \leq PL_o(I)$, is a countable union of pairwise disjoint open intervals in $(0, 1)$. Let the collection $\mathcal{O}_H$ always denote the countable set of open, pairwise disjoint intervals of the support of H . We call these intervals the *orbitals* of H . There is a natural total order on $\mathcal{O}_H$, where if $A, B \in \mathcal{O}_H$, where $A \neq B$, we will say $A < B$ or A is to the left of B if and only if given any $x \in A$ and $y \in B$, we have $x < y$ under the natural order induced by $I \subset \mathbf{R}$. Since A and B are disjoint, connected subsets of $\mathbf{R}$, this definition does not depend on the choices of x and y .

If the collection $\mathcal{O}_H$ is finite, we may speak of the “first” orbital, or “second” orbital, etc., where the first orbital is the leftmost orbital under the definition given above, the second orbital is the orbital to the left of all other orbitals in $\mathcal{O}_H$ other than itself and the first orbital, and so on.

Given an open interval $A = (a, b) \subset \mathbf{R}$, where $a < b$, we will refer to a as the *leading end* of A , and to b as the *trailing end* of A . If the interval is an orbital of some group $H \in PL_o(I)$, we will refer to the *ends of the orbital* in the same fashion.

If $h \in H$ and $x \in \text{Supp}(h)$, we will say that h moves x to the left if $xh < x$, and we will say that h moves x to the right if $xh > x$. Furthermore, we will say that $x \in I$ is a *breakpoint* for h if the left and right derivatives of h exist at x , but are not equal. We recall that by definition, h will admit only finitely many breakpoints. If $\mathcal{B}_h$ represents the set of breakpoints of the element h , then $(0, 1) \setminus \mathcal{B}_h$ is a finite collection of open intervals, which we will call *affine components* of h , which admit a natural “left to right” ordering as before. We shall unhesitatingly refer to the “first” affine component of h , or the “second” affine component of h , etc. We sometimes will refer to the first affine component of h as the *leading affine component* of h , and to the last affine component of the domain of h as the *trailing affine component* of h .

Given an element $h \in H$ the group $\langle h \rangle$ generated by h has its own orbitals, which we will typically call the *orbitals* of h . Any such collection of orbitals for an element $h \in PL_o(I)$ is finite by Remark 2.5, and we will denote the number of orbitals for such an element h as o_h . We will denote the ordered (left to right, as before) collection of orbitals of h as $\mathcal{A}^h = \{A_i^h\}_{i=1}^{o_h}$.

The following are some useful remarks.

- Remark 2.5**
1. If A is an orbital for $h \in H$, then either $xh > x$ for all points x in A , or $xh < x$ for all points x in A .
 2. Any element $h \in PL_o(I)$ has only finitely many orbitals.
 3. If $h \in PL_o(I)$ and $A = (a, b)$ is an orbital of h , then given any $\epsilon > 0$ and x in A , there is an integer n so that $|xh^{-n} - a| < \epsilon$ and $|xh^n - b| < \epsilon$.

pf:

For the first point, the difference of x and xh can never be zero in an orbital, but the difference function is continuous, so the intermediate value theorem of calculus implies the difference is always positive, or always negative, throughout the orbital.

For the second point, assume it is incorrect, then, h will behave as the identity function for an infinite sequence of points $(x_i)_{i=1}^\infty$ in I so that between any two points x_i and x_{i+1}

there would be a point y_i so that $y_i h \neq y_i$. Hence, h would have to admit to infinitely many affine components, which contradicts the definition of $PL_o(I)$.

For the final point, suppose that h moves points to the right on its orbital A (if not, a symmetric argument will prove the result), and let $\epsilon > 0$.

The sequence $y_i = xh^i$ defined for all natural numbers i is an increasing sequence of points which is bounded above by b . In particular, it must have a limit y (greater than all of the x_i) by the completeness of the real numbers. But now $yh = y$ by the continuity of h , so y must be b . Hence, there is a natural number m_1 so that $|y_{m_1} - b| = |xh^{m_1} - b| < \epsilon$. A symmetric argument shows that there is a natural number m_2 so that $|xh^{-m_2} - a| < \epsilon$. Let $n = \max(m_1, m_2)$. $\diamond$

Given an orbital A of H we say that h *realizes an end of A* if some orbital of h lies entirely in A and shares an end with A . Note that Brin uses the word “Approaches” for this concept in [1], but we will use “Approaches” to also indicate the direction in which h moves points, as follows: we will say that h *approaches the end a of A in A* if h has an orbital B where $B \subset A$ and B has end a , and h moves points in B towards a . In particular, h realizes a in A and h moves points in its relevant orbital towards a . If A is an orbital for H then we say that $h \in H$ *realizes A* if A is also an orbital for h .

If g and h are elements of $PL_o(I)$ and there is an interval $A = (a, b) \subset I$ so that both g and h have A as an orbital, then we will say that g and h *share the orbital A* .

Given $g \in PL_o(I)$, and any $x \in (0, 1)$, there is an $\epsilon > 0$ so that $(x, x + \epsilon)$ is contained in an affine component C_x of h . The slope of h on C_x is the value of the right hand derivative $h'_+(x)$ of h at x . Given another element $h \in PL_o(I)$, we can find the maximal subinterval $[0, a] \subset [0, 1]$ so that for any $x \in [0, a]$ we have $h(x) = g(x)$. If $a = 1$, then $h = g$. If $a < 1$, then $h'_+(a) \neq g'_+(a)$. In particular, we can define a total order on the set of elements of $PL_o(I)$, by defining $g < h$ if $g'_+(a) < h'_+(a)$, where a is the largest point in $[0, 1]$ where g and h are identical on $[0, a]$. We will call this the *left total order on $PL_o(I)$* .

2.4.1 Conjugation and orbitals

In this section, we will establish some notation to help us to more flexibly pass to subgroups of a group in $PL_o(I)$. This section will help us to understand the orbital structure of conjugates of an element in $PL_o(I)$.

Let $g, h \in PL_o(I)$ and let $k = g^h = h^{-1}gh$. Suppose that $\mathcal{O}_g = \{A_i\}_{i=1}^n$ are the $n = o_g$ orbitals of g in left to right order, where $n \in \mathbf{N}$ and $i \in \{1, 2, \dots, n\}$. Define

$$A_i^* = \{x \in I | xh^{-1} \in A_i\} = A_i h$$

for all $i \in \{1, 2, \dots, n\}$.

The following is standard.

Lemma 2.6 $o_k = o_g = n$ and collection $\{A_i^*\}_{i=1}^n$ is the ordered set of orbitals of g^h in left to right order.

pf: Throughout this proof, let i always refer to an index with $1 \leq i \leq n$.

Suppose $x \in I$ so that xh^{-1} is not an element in any A_i . Then $xg^h = xh^{-1}gh = xh^{-1}h = x$, since xh^{-1} is in the fixed set of g . In particular, the support of g^h is in the union of the

sets A_i^* . Now suppose $x \in A_i^*$ for some index i . Then $xh^{-1} \in A_i$, now $xg^h = (xh^{-1}g) \cdot h \neq x$, since g moves the point xh^{-1} . In particular, each A_i^* is contained in the support of g^h . Now we see that the support of g^h is precisely the union of the finite collection of sets A_i^* . Now, the collection of A_i^* 's is a finite collection of pairwise disjoint, open intervals, being the homeomorphic image of a finite collection of pairwise disjoint, open intervals. In particular, each A_i^* is a connected component of the support of g^h , and is therefore an orbital of g^h . Since the set of orbitals $\{A_i\}_{i=1}^n$ is ordered in left to right fashion for g , and since h is order preserving, the indexed collection $\{A_i^*\}_{i=1}^n$ of the orbitals of g^h is also ordered in left to right fashion. $\diamond$

In the setting of the above lemma, we will say that the $A_i h$ are the *induced orbitals* of k from g by the action of h . We might also say that the orbitals of k are induced from the orbitals of g by the action of h .

The following two unrelated points are worth pointing out:

- Remark 2.7** 1. Suppose $g, h \in PL_o(I)$ and $f = gh$. If b is a breakpoint of f then b is a breakpoint of g or bg is a breakpoint of h .
2. Let $\alpha, \beta \in PL_o(I)$, then α^β has the same leading and trailing slopes on its orbitals as α has on each of its corresponding orbitals.

pf:

For the first point, suppose b is a breakpoint of f , but b is not a breakpoint of g . The lefthand slope of f at b is the product of the lefthand slopes of g at b and h at bg . Also, the righthand slope of f at b is the product of the righthand slopes of g at b and h at bg . Since the lefthand and righthand slopes of g at b are the same, the lefthand and righthand slopes of g at bh must be different. In particular, bg is a breakpoint for h .

For the second point, let $A = (a, b)$ be an orbital of α^β , corresponding to an orbital $A' = (a', b')$ of α . There is $c \in (a, b)$ so that (a, c) is contained in an affine component of β^{-1} , so that $(a', c') = (a, c)\beta^{-1}$ is in an affine component of α , and so that $(a', c'') = (a', c')\alpha$ is contained in an affine component of β . By Remark 2.7, (a, c) is therefore contained in an affine component of α^β , and hence the slope of α^β on (a, c) is the leading slope of α^β on A , since the leading affine component of α^β with non-trivial intersection with A must now contain (a, c) . The slope of α^β on (a, c) is the product of the slope of β^{-1} on (a, c) , the slope of α on (a', c') , and the slope of β on (a', c'') . But the functions β and β^{-1} are inverse, so if x is in an affine component of β^{-1} with slope s , then $y = x\beta$ is in an affine component of β with slope $1/s$. Now then the image of (a, c) under β^{-1} must be in an affine component (a', d) of β , and the slopes of β^{-1} on (a, c) and β on (a', d) are multiplicative inverses. But now we see that $(a', c'') \subset (a', d)$, so in particular, the lead slope of α^β on (a, b) is the lead slope of α on (a', b') .

The argument for the trailing slopes is similar. $\diamond$

2.4.2 Signed orbitals

We give this discussion its own section, since the definitions we discuss here are new, and are of extreme importance to understanding the arguments to follow. Otherwise, this could easily be included at the end of the previous section.

Let $1 \neq g \in G$ and A be an orbital of g . It is immediate that A is realized by infinitely many elements of G (for instance, non-zero powers of g will always realize A). Often, we want to associate some specific element of G which realizes the orbital A with A .

Given a group $G \leq PL_o(I)$, we define a *signed orbital* (in G) to be a pair (A, g) where A is an orbital of the element $g \in G$.

Now we will build up some convenient language around signed orbitals, and note some trivial facts. Given $\gamma = (A, g)$, a signed orbital of G , we will call A the *orbital of γ* , and g the *signature of γ* . We observe, as before, that if n is a non-zero integer then (A, g^n) is another signed orbital with orbital A . In particular, if (A, g) is a signed orbital, then so is (A, g^{-1}) . Note in this case that if g moves points to the right on A , then g^{-1} moves points to the left on A , and vice-versa, so that we will often replace a signed orbital by the signed orbital consisting of the same orbital, but with the inverse of the original signature, so that the new signature moves points in a desired direction on the orbital.

Given a set P of signed orbitals, we will often refer to two induced sets from P ; the set of orbitals of P , denoted O_P , and the set of signatures of P , denoted S_P . For completeness,

$$\begin{aligned} O_P &= \{A \subset [0, 1] \mid \text{there is } g \in G \text{ with } (A, g) \in P\}, \\ S_P &= \{g \in G \mid \text{there is } A \text{ an orbital of } g \text{ with } (A, g) \in P\}. \end{aligned}$$

Noting the standard (left) total order on elements of $PL_o(I)$, we observe that the set of signed orbitals of $PL_o(I)$ is partially ordered by using the lexicographical ordering, where we use the partial order on subsets of I (induced via inclusion) for the first coordinate, and the left total order on elements of $PL_o(I)$ for the second coordinate. In particular, the set of signed orbitals of any subgroup of $PL_o(I)$ is also a poset.

Suppose H is a subgroup in $PL_o(I)$, and H has two elements α and β , which then generate some subgroup G of H . If we restrict our attention to the orbitals of G , one orbital at a time, we discover that the orbitals of α and β must be arranged so that the orbitals of α cover the fixed set of β within each orbital of G , and vice-versa. This arrangement of orbitals allows us to move points from one side of an orbital of G to the other, by repeated application of the elements α and β , and of their inverses, at the appropriate moments. A detailed argument that we can move any point inside an orbital A of H arbitrarily close to either end of A will be given shortly.

Let $\mathcal{C} = \{O_i\}_{i=1}^n$ be a set of signed orbitals of G , so that $O_i = (A_i, g_i)$ for each $i \in \{1, \dots, n\}$. Suppose that $\mathcal{C}$ satisfies the following properities.

1. For any $k \in \{1, \dots, n-1\}$, $A_k \cap A_{k+1} \neq \emptyset$.
2. For any $k \in \{1, \dots, n\}$, there is a point $p_k \in A_k$, which is not in any A_j for $j \neq k$.
3. For any $k \in \{1, \dots, n-1\}$, $p_k < p_{k+1}$.

then we will call $\mathcal{C}$ a *transition chain for the group G* . The integer n will be called the *length of the transition chain $\mathcal{C}$* .

Given a transition chain $\mathcal{C}$ of length n , $\mathcal{C} = \{(A_i, g_i) \mid i \in \{1, \dots, n\}\}$, we fix the notation $A_{\mathcal{C}} = \bigcup_{i=1}^n A_i$, as it will often be convenient to refer to the set $A_{\mathcal{C}}$.

If $\mathcal{C}$ is a transition chain with length n as above, and $1 \leq j \leq k \leq n$, then $\mathcal{C}' = \{O_i\}_{i=j}^k$ will be called a *contiguous subset of $\mathcal{C}$* . Then it is an immediate observation that any

contiguous subset of a transition chain is also a transition chain. Given two transition chains $\mathcal{C}'$ and $\mathcal{C}$, we will say $\mathcal{C}' \leq \mathcal{C}$ whenever $\mathcal{C}'$ is a contiguous subset of $\mathcal{C}$. Given a group G , its set of transition chains forms a partially ordered set under this binary relation.

Here is a useful lemma:

Lemma 2.8 *If $H \leq PL_o(I)$ and $A = (a, b)$ is an orbital for H , then given any points $c, d \in A$, with $c < d$, there is an element $g \in H$ so that $cg > d$.*

pf: $[c, d]$ is contained in an orbital of H , and therefore it is contained in the union of the orbitals of the elements of H . Since $[c, d]$ is compact, it follows that it is covered by a finite subcollection $\mathcal{C}'$ of the orbitals of the elements of H . This implies that there is a smallest positive integer n with a transition chain $\mathcal{C} = \{(A_i, g_i) | 1 \leq i \leq n, i \in \mathbf{N}\}$ for H whose orbitals cover $[c, d]$. If $n = 1$ we are done by Remark 2.5, so assume $n > 1$. We note in passing that c is an element of A_1 but not of A_2 and d is an element of A_n but d is not an element of A_{n-1} .

Improve $\mathcal{C}$ by supposing we chose signatures intelligently, so that each $(A_i, g_i) \in \mathcal{C}$ satisfies the property that g_i moves points to the right on A_i .

Now, by the definition of transition chain, we know that $B_i = A_i \cap A_{i+1}$ is actually a non-empty open interval for each integer i where $1 \leq i < n$. Let δ_1 be the minimum length of these intervals B_i . By the definition of transition chain, each orbital A_i has length at least δ_1 (and $2\delta_1$ for interior orbitals in the chain), since each such orbital has a distinguished point which is outside of the intersection of this orbital with the other orbitals of the transition chain. Let δ_2 be the minimum of the two distances, one from the left end of A_1 to c , and the other from d to the right end of A_n . Now let $\delta = \min(\delta_1, \delta_2)/2$. It is immediate by construction that each orbital of $\mathcal{C}$ has length greater than 2δ , and also that each interval B_i created above has length greater than 2δ . For each $k \in \{1, 2, \dots, n\}$, let x_k be a point in A_k a distance less than δ from the left side of A_k . By Remark 2.5, for each integer k where $1 \leq k \leq n$, there is a positive integer m_k so that $g_k^{m_k}$ will take x_k to a point y_k within δ of the right end of A_k . Observing that for all integers k where $1 \leq k < n$ we have that $y_k > x_{k+1}$, we see that $g = g_1^{m_1} g_2^{m_2} \cdots g_n^{m_n}$ moves x_1 to the right of y_n . But now $x_1 < c < d < y_n$ by construction, so $cg > d$. $\diamond$

2.5 Thompson's group F and balanced subgroups of $PL_o(I)$

Brin showed in [1] the following theorem:

Theorem 2.9 (Ubiquitous F) *If a group $H \leq PL_o(I)$ has an orbital A so that some element $h \in H$ realizes one end of A , but not the other, then H will contain a subgroup isomorphic to Thompson's group F .*

The condition on the orbital is weak enough that one readily observes that “ $PL_o(I)$ is riddled with copies of F ”, quoting Brin. Hence it becomes a natural question to ask what can be said about subgroups of $PL_o(I)$ which have no orbitals satisfying the ubiquity condition.

We will say that an orbital A of a group $H \leq PL_o(I)$ is *imbalanced* if some element $h \in H$ realizes one end of A , but not the other, and we will say A is *balanced* if whenever an element $h \in H$ realizes one end of A , then h also realizes the other end of A (note that

h might do this with two distinct orbitals). Extrapolating, given a group $H \leq PL_o(I)$, we will say that H is *balanced* if given any subgroup $G \leq H$, and any orbital A of G , every element of G which realizes one end of A also realizes the other end of A . Informally, H has no subgroup G which has an orbital that is “heavy” on one side. In the case where H has a subgroup G with an imbalanced orbital, then we will say that H is *imbalanced*.

Remark 2.10 *If $H \leq PL_o(I)$ and H is imbalanced, then H has a subgroup isomorphic to Thompson’s group F .*

Since F' is non-trivial and simple ([4], Theorem 4.3), F is not solvable. Thus imbalanced groups are not solvable.

Possibly less obvious is that the dynamics of balanced groups are much easier to understand than those of imbalanced groups. We will trade heavily on this in the remainder, so the next few subsections will establish some of the common tools that we will have available to us when we are analyzing balanced groups. In the meantime, we build some tools to help us “find” imbalanced groups.

2.5.1 A useful homomorphism

Let us suppose that $H \leq PL_o(I)$ and $A = (a, b)$ is an orbital of H . To simplify the arguments for now, let us suppose that A is the only orbital of H . We can define a map $\phi : H \rightarrow \mathbf{R} \times \mathbf{R}$ defined by $h \mapsto (h_a, h_b)$ where $h_a = \ln(h'_+(x))$ and $h_b = \ln(h'_-(x))$. Ie., we take the logs of the slopes of h at the ends of A . Since h is a p.l. orientation-preserving homeomorphism of I , we see that the derivatives exist and are positive, and so ϕ is well defined for all $h \in H$. If h does not realize a (resp. b) then we see that h behaves as the identity near a (b) in A , and so $h_a = \ln(1) = 0$ ($h_b = 0$). If $h, g \in H$ then $hg\phi = h\phi + g\phi$ in $\mathbf{R} \times \mathbf{R}$ by the chain rule. In particular, we see the following remark:

Remark 2.11 *ϕ is a homomorphism of groups.*

Now the image of ϕ is quite interesting, it carries a small amount of the complexity of H , but still enough to allow us to find out if A is an imbalanced orbital.

Lemma 2.12 *The orbital A is imbalanced if and only if $Im(\phi)$ contains an element of the form $(\alpha, 0)$ or $(0, \alpha)$ where $\alpha \neq 0$.*

pf: Suppose that $h \in H$ approaches one end of A but not the other. Then the slope of h near one end is 1, but on the other end is not 1. In particular, $h\phi = (h_a, h_b)$ is either $(0, \alpha)$ or $(\alpha, 0)$ for some non-zero α . Hence, if A is imbalanced, then $Im(\phi)$ contains an element in this form. Conversely, if $Im(\phi)$ has an element r_α of the form $(0, \alpha)$ or $(\alpha, 0)$, where α is not zero, then there is an $h \in H$ so that $h\phi = r_\alpha$. For this h , either $h_a = 0$ and $h_b = \alpha$, in which case h realizes b but does not realize a , or, in the other case, $h_a = \alpha$ and $h_b = 0$, where we see that h realizes a but not b . In either case, we see that A is an imbalanced orbital. $\diamond$

This next technical lemma will help with the lemma that follows it:

Technical Lemma 2.13 *Suppose $H \leq PL_o(I)$ and H has an orbital $A = (a, b)$, and that H has a sequence of elements $(g_n)_{n=1}^\infty$ in H which satisfies the properties below.*

1. For each $i \in \mathbf{N}$, the lead slope of g_{i+1} in A is less than the lead slope of g_i in A .
2. Given any real number $q > 1$, there is an $i \in \mathbf{N}$ so that the lead slope of g_i in A is p where $1 < p < q$.

Then there is $c \in (a, b)$ so that given any real number $s > 1$, H has an element α which has an affine component Γ containing (a, c) , and with slope r on Γ where $1 < r < s$.

pf:

To simplify the language of this argument, we will restrict our attention to the orbital A , treating it as the domain of the elements of H , so that the phrase “The first affine component of $[h \in H]$ ” will really mean the open interval (a, u) where h has an affine component of the form (v, u) where $v \leq a$. We will also refer to this as the “leading (or lead) affine component of h ”. Similarly, we will refer to the slope of h on its leading affine component as the “lead slope of h ”.

Note that the second condition on $(g_n)_{n=1}^\infty$ implies that every element of $(g_n)_{n=1}^\infty$ has lead slope greater than one.

For each $i \in \mathbf{N}$, let (a, b_i) be the first affine component of g_i , and let s_i represent the lead slope of g_i .

We are now in a position to define a new sequence of functions $(h_i)_{i=1}^\infty$ which satisfy the following conditions:

1. For each $i \in \mathbf{N}$, the lead slope of h_i is s_i .
2. For each $i \in \mathbf{N}$, the leading affine component of h_i contains (a, b_1) .

At this point, by taking $c = b_1$, we see that given any $s > 1$, by the hypotheses on the g_i there will be an $N \in \mathbf{N}$ so that for all $n > N$ we will have that $s > s_n > 1$, and that h_n has leading affine component containing (a, c) .

For each $i \in \mathbf{N}$, define n_i to be the smallest non-negative integer so that $b_1 g_1^{-n_i} < b_i$, and define $h_i = g_1^{-n_i} g_i g_1^{n_i}$. Since g_1 moves points to the right on its first affine component, n_i is well defined, and therefore h_i is well defined. We now check that h_i satisfies the two conditions, for each i . Firstly, we observe that the lead slope of each h_i is the product of the lead slopes of the elements of the product $g_1^{-n_i} g_i g_1^{n_i}$, which is $(\frac{1}{s_1})^{n_i} s_i s_1^{n_i} = s_i$. Secondly, by Remark 2.7, we know that if $x \in (a, b)$ is a breakpoint of the product $g_1^{-n_1} g_1 g_1^{n_1}$, then the image of x under application of some initial partial product (possibly empty) must be a breakpoint of the next term of the overall product. However, the first breakpoint of g_1^{-1} is $d_1 = b_1 g_1 > b_1$, and g_1^{-1} moves points left in its first affine component so if $x \in (a, b_1)$, then $x g_1^{-k} < d_1$ for all non-negative integers k . In particular, if $x \in (a, b_1)$, then x is in the first affine component of $g_1^{-n_i}$ for any $i \in \mathbf{N}$. Given an $i \in \mathbf{N}$, the first breakpoint of g_i in A is b_i , but n_i was chosen so that $b_1 g_1^{-n_i} < b_i$, so if $x \in (a, b_1)$, then the image of x under $g_1^{-n_i}$ is in the first affine component of g_i , in particular, (a, b_1) is contained in the first affine component of $g_1^{-n_i} g_i$. Finally, the first breakpoint of $g_1^{n_1}$ is the image of b_1 under $g_1^{-n_1+1}$, but $b_1 g_1^{-n_i} g_1 \geq b_1 g_1^{-n_i} g_i$ because the leading slope of g_i is less or equal to the leading slope of g_1 . In particular, the whole interval (a, b_1) is in the first affine component of h_i . $\diamond$

The following lemma is the achievement of the section, as it will enable us to find the remarkable “controllers”; elements that control the global behavior of a balanced group on its orbitals.

Lemma 2.14 *Suppose H is a balanced group with single orbital $A = (a, b)$, and ϕ is the log-slope homomorphism defined before, then $\phi(H) \cong \mathbb{Z}$ or $\phi(H)$ is trivial in $\mathbf{R} \times \mathbf{R}$.*

pf: Let H be a balanced subgroup of $PL_o(I)$. Each element of H either realizes both ends of A , or neither. In particular, the group homomorphism $\rho_1 : \mathbf{R} \times \mathbf{R} \rightarrow \mathbf{R}$ which is projection on the first factor has the property that $\ker(\rho_1) \cap \text{Im}(\phi) = \{(0, 0)\}$, the trivial subgroup of $\mathbf{R} \times \mathbf{R}$. This implies that the image of ϕ in $\mathbf{R} \times \mathbf{R}$ is isomorphic to the group $H_1 \leq \mathbf{R}$ obtained by considering only the first factors of elements of $\text{Im}(\phi)$. If no element realizes the ends of A , then $H_1 = \{0\}$, the trivial (additive) group, and we are done. Therefore, let us suppose instead that some elements in H realize the left end of A (and therefore also the right) so that H_1 cannot be the trivial subgroup of $\mathbf{R}$.

If H_1 is discrete in $\mathbf{R}$ then H_1 is either trivial, or isomorphic to $\mathbf{Z}$, but by assumption, H_1 is not the trivial group, hence in this case $H_1 \cong \mathbf{Z}$. Hence, we shall suppose that H_1 is not discrete in $\mathbf{R}$. In this case, by taking the difference of two elements in the image which are very near each other, we see that we can find an element of H_1 which is as close to zero as we like. This implies there are elements of H whose leading slopes are as close to one as we like, without actually being one. If h is an element with leading slope $s \neq 1$, then one of h or h^{-1} has slope greater than one, since the leading slope of h is s , but the leading slope of h^{-1} is $1/s$ (note that s cannot be zero, since no element of $PL_o(I)$ has an affine component with slope zero).

Now suppose that H is abelian. By a result of Brin and Squier [3], if two elements in $PL_o(I)$ commute and share a common orbital, then their projections on that orbital have a common root. If two elements have non-disjoint support and commute, it is easy to see that the intersections of their supports actually is a set of commonly shared orbitals. Now, let h be some element of H with leading slope $s > 1$. For each positive integer n , let g_n be an element of H with leading slope s'_n where $1 < s'_n < \frac{n+1}{n}$. Now for each g_n , the pair h and g_n has a common root h_n (on their leading orbital), but infinitely many of the roots h_n have pairwise distinct leading slopes, since these slopes are always less than or equal to the slopes of the g_n , and in particular, h must then have infinitely many distinct roots in H on its leading orbital. By another result in [3], no element of $PL_o(I)$ has infinitely many distinct roots, so we must conclude that H is not abelian.

Note that by the details of the previous paragraph, it is easy to construct a countably infinite sequence of elements $(g_n)_{n=1}^\infty$ in H which satisfies the properties below:

1. For each $i \in \mathbf{N}$, the lead slope of g_{i+1} is less than the lead slope of g_i .
2. Given any real number $r > 1$, there is an $i \in \mathbf{N}$ so that the lead slope of g_i is s where $1 < s < r$.
3. given $i, j \in \mathbf{N}$, we will have $[g_i, g_j] = 1$ implies $i = j$.

Therefore, by Lemma 2.13 there is $c \in (a, b)$ and elements of H with lead slopes that are greater than but arbitrarily close to one, and whose leading affine components contain (a, c) .

Let f and g be two elements of H with $h = [f, g] \neq 1$. The fixed set of h in I is disconnected, and contains two components of the form $[0, u']$ and $[v', 1]$ for some numbers $u', v' \in (a, b)$. In particular $\inf(\text{Supp}(h)) = u'$ and $\sup(\text{Supp}(h)) = v'$.

By Lemma 2.8 there is $\alpha \in H$ so that $v'\alpha < c$, so that $j = h^\alpha$ has all of its orbitals inside (a, c) . Let $u = \inf(\text{Supp}(j))$ and $v = \sup(\text{Supp}(j))$. In particular, $v < c$.

We will now perturb j slightly via a conjugation which will move the orbitals of j to the right by a distance less than L , so that j and the new element together will generate a group with an imbalanced orbital.

Suppose $L > 0$ is smaller than two particular lengths. The first length is the length of the second component of the fixed set of j in A which has non-zero length (note, this component might just be $[v, b)$, if j has only two such), and the second length is the length of the first orbital of j .

Choose an element $\beta \in H$ whose leading affine component in A contains (a, c) and whose lead slope is greater than one, but so near one that no point in (a, c) will move to the right a distance greater than L . Now the elements j and j^β will generate a group G with leading orbital (u, w) where j realizes u and possibly w (if the right ends of the appropriate orbitals of j and j^β are aligned), while j^β will achieve w but not u . To see this, note that the left end of the first orbital of j^β is in the first orbital of j , so that u is the left end of the leading orbital of G , and only j realizes it. Meanwhile, the right end of the first orbital of j^β is to the right of the right end of the first orbital of j , so that the leading orbital of j^β contains the right end of the leading orbital of j . As we progress to the right, if the solitary fixed point sets of j and j^β align before we reach the second component of the fixed set of j in A with non-zero length, then the right end of the first orbital of G will be achieved by both j and j^β . Otherwise, the first orbital of G will extend rightward into the interior of the second fixed set of j with a non-empty interior, so that the right end of this orbital of G will be realized only by j^β . In all cases, G will be imbalanced, and hence H will also be imbalanced. This contradicts the hypotheses of the lemma, and so we see that H_1 must be discrete in $\mathbf{R}$, and therefore the image of ϕ is isomorphic to $\mathbf{Z}$ or the trivial group in $\mathbf{R} \times \mathbf{R}$. $\diamond$

The kernel of the homomorphism ϕ is naturally very important as well, it is the subgroup of H which consists of elements which are the identity near the ends of A . Typically, we will refer to this normal subgroup as $\hat{H}$.

2.5.2 Controllers

A consequence of section 2.5.1 is that the structure of a balanced group with one orbital is very special. In this section we will explore this idea further.

Lemma 2.15 (Balanced Generator Existence) *Suppose that H is a balanced subgroup of $PL_o(I)$ with single orbital A , that there is some element in H which realizes an end of A , and that $\hat{H}$ is the subgroup of H which consists of all elements in H which are the identity near the ends of A . Then there is an element g of H so that $H = \langle \langle g \rangle, \hat{H} \rangle$, where g realizes both ends of A .*

pf:

Let Γ_A be the set of elements of H which realize both ends of A . By our assumptions, Γ_A is not empty. Now observe that $H = \langle \langle \Gamma_A \rangle, \hat{H} \rangle$.

By lemma 2.14 the image $\phi(H)$ is cyclic in $\mathbf{R} \times \mathbf{R}$. Let γ be a generator of the image of ϕ . Let g be an element of H so that $g\phi = \gamma$. We observe that since γ is non-trivial in both

components, g realizes both ends of A . Since γ generates the image of ϕ , if $\hat{g} \in \Gamma_A$, then $\hat{g}\phi = g^k\phi$ for some $k \in \mathbf{Z}$. Hence, $g^{-k} \cdot \hat{g} \in \hat{H}$. This now implies that $\Gamma_A \subset \langle g, \hat{H} \rangle$, so that $\langle g, \hat{H} \rangle = \langle \Gamma_A, \hat{H} \rangle = H$. $\diamond$

We will call any element c of a balanced group H with one orbital A , which satisfies the rule $H = \langle \langle c \rangle, \hat{H} \rangle$, a *controller* of H . A controller of H is clearly a special element.

Given a controller c of a balanced group H with one orbital A , we can write any element h of H uniquely in the form $c^k \cdot \overset{\circ}{g}$, where k is some integer, and $\overset{\circ}{g} \in \hat{H}$. We will call this the *c-form* of h .

We will say that a controller c of the group H is *consistent* if and only if its image $(\alpha, \beta) = c\phi$ satisfies the property that $\text{sign}(\alpha) = -\text{sign}(\beta)$. Otherwise we will say the controller is *inconsistent*. The idea behind this definition is that a one bump controller should be consistent, since it is either moving points to the right everywhere on its support, or it is moving points to the left everywhere on its support. An inconsistent controller must have a fixed point set, and has at least one bump where the controller moves points to the right, and one bump where the controller moves points to the left. It turns out that a consistent controller actually is a one-bump element of H .

Lemma 2.16 *Suppose H is a balanced subgroup of $PL_o(I)$ and H has unique orbital A . Further suppose that H has a consistent controller c , then c realizes A .*

pf:

Since c and c^{-1} are both controllers, and either both satisfy or both fail the conclusion of the statement of the lemma, we will assume c moves points to the right on its orbitals near the ends of A . Suppose c has a non-trivial fixed set K in A . K is closed and bounded and hence compact. Let $u = \inf K$ and $v = \sup K$, so that (a, u) is the first orbital of c and (v, b) is the last orbital of c . Now there are points $x \in (a, u)$, and $y' \in (v, b)$ so that we have $a < x < u \leq v < y' < b$. By Lemma 2.8 there is an element $g \in H$ so that $xg > y'$. Writing g in its c -form, we have that $g = c^k \overset{\circ}{g}$ for some integer k and element $\overset{\circ}{g} \in \hat{H}$. In particular, the element $h = gc^{-k}$ is trivial near the ends of A , but still satisfies $xh = y > v$. The element h therefore has an orbital $D = (r, s)$ which spans the fixed set K of c . Suppose $e = \inf(\text{Supp}(h))$, so that $a < e \leq r$. By Lemma 2.5 there is a positive integer N_1 so that for any integer $n_1 > N_1$ we will have $r < ec^{n_1} < u$. Suppose $f = \sup(\text{Supp}(h))$, so that $s \leq f < b$. By Lemma 2.5 there is a positive integer N_2 so that for any integer $n_2 > N_2$ we will have $f < sc^{n_2} < b$. Let $n = \max(N_1, N_2)$, then, the element $j = h^{(c^n)}$ has its first orbital starting at some interior point of (r, s) , and its orbital induced from (r, s) has right end t which is strictly to the right of f . In particular the group $H_1 = \langle j, h \rangle$ has an orbital $B = (r, t)$, where we note that $t > s$ by construction. Now, h realizes the left end of B in B , but not the right, hence H_1 , and therefore H , is imbalanced. But this contradicts our assumptions, therefore c must have orbital A . $\diamond$

Corollary 2.17 *Suppose H is a balanced subgroup of $PL_o(I)$ and H has a unique orbital $A = (a, b)$. If H has a consistent controller, c , then any element g of H which realizes both ends of A actually realizes A .*

pf: Let $g = c^k \overset{\circ}{g}$ be the c -form of g , and suppose that g has a non-empty fixed set in (a, b) . The behavior of g near the ends of A (beyond the support of $\overset{\circ}{g}$) depends entirely on c

and the integer k , and therefore g is either moving points to the right on both of its leading and trailing orbitals, or moving points to the left on both of its leading and trailing orbitals, since k cannot be zero. Now the details of the argument of Lemma 2.16 show that H must be imbalanced, which contradicts our assumptions, hence g must have orbital A . $\diamond$

We will now consider the case where H has an inconsistent controller.

Remark 2.18 *Suppose H is a balanced subgroup of $PL_o(I)$ and H has unique orbital $A = (a, b)$. If H has an inconsistent controller c , then no element of H realizes A .*

pf: Any element h which realizes both ends must do so inconsistently. Hence the difference function $D : [a, b] \rightarrow \mathbf{R}$ defined by the rule $xD = xh - x$ is a continuous function which is positive near one end of A and negative near the other, and so by the intermediate value theorem, there is some point $x \in A$ so that $xD = 0$. The point x corresponds to a fixed point of h . $\diamond$

In the case where H is a one-orbital balanced group with an inconsistent controller, by Remark 2.18 any element h which realizes both ends of A must have a non-trivial fixed set F_h in A . Suppose there exists an element $\overset{\circ}{g}$ of $\hat{H}$ with an orbital $A_{\overset{\circ}{g}}$, where the ends of $A_{\overset{\circ}{g}}$ are in differing orbitals of h . The element $\overset{\circ}{g}$ enables the transfer of a point $x \in \text{Supp}(h) \cap A_{\overset{\circ}{g}}$ from its own orbital in h to another orbital of h . In particular, the point xg^k will be in a different orbital of h than the orbital which contains x , for some integer power k . Such an element $\overset{\circ}{g}$ is very useful to have in hand, so we will study questions about the existence and structure of such elements in a one-orbital balanced group with inconsistent controller.

Lemma 2.19 *Suppose H is a balanced subgroup of $PL_o(I)$ and H has a unique orbital $A = (a, b)$. Suppose that H has an inconsistent controller c and let $\hat{H}$ represent the normal subgroup of H consisting of the elements in H which achieve neither end of A . Now, the fixed set $F_c = \text{fix}(c) \cap A$ of c in A is contained in an orbital of some element $\overset{\circ}{g} \in \hat{H}$.*

pf: Let $x = \inf F_c$ and let $y = \sup F_c$. Note that $x, y \in F_c$. There is an element $g \in H$ so that $xg > y$ by lemma 2.8. We can write $g = c^k \cdot \overset{\circ}{g}$ for some integer k and some element $\overset{\circ}{g} \in \hat{H}$. Clearly, c^{-k} does not move the fixed set of c , but g moves a closed interval containing the fixed set of c completely off of itself. In particular, we have that $xc^{-k}g > y$, so that $F_c \subset [x, y] \subset \text{Supp}(c^{-k}g)$. But $c^{-k}g = \overset{\circ}{g} \in \hat{H}$, so $F_c \subset B$ for some orbital B of $\overset{\circ}{g}$. $\diamond$

Corollary 2.20 (Transfer Existence) *Suppose H is a balanced subgroup of $PL_o(I)$ and H has a unique orbital $A = (a, b)$. Suppose that H has an inconsistent controller c and let K be any compact set in (a, b) . Then K is contained in an orbital of some element $\overset{\circ}{g} \in \hat{H}$.*

pf:

Suppose that c moves points to the left on its first orbital, and moves points to the right on its last, by replacing c with its inverse, if necessary.

Let $x = \inf K$ and $y = \sup K$. Let $\hat{g} \in \hat{H}$ be an element which contains the fixed set of c in some orbital $\hat{A}_c$. By the constraints that c moves points to the left on its first orbital, that c moves points to the right on its last orbital, and that the conjugate $c^{-1}\hat{g}c$ has the induced orbital corresponding to $\hat{A}_c$ larger than $\hat{A}_c$. By conjugating repeatedly with c , we can make

the ends of this orbital approach the ends of A , so that after some conjugation the resulting orbital will contain the set $[x, y]$. The resulting function is $\overset{\circ}{g} \in \tilde{H}$. $\diamond$

Suppose that we know that A is an orbital of a balanced group H , and that the ends of A are achieved by some element g in H . Let H_A be the set of elements of $PL_o(I)$ each of which is equal to the restriction of some element of H on the orbital A , and behaves as the identity off of A . H_A is trivially a group with unique orbital A , and is a quotient of H . We will call H_A the projection of H on A .

We will now generalize our language somewhat. Let H be a subgroup of $PL_o(I)$ with an orbital A , and let H_A be the projection of H on A . H_A has a controller $\tilde{c}$ for A . Let $\rho : H \rightarrow H_A$ be the projection homomorphism on the orbital A . Let $T = \{c \in H \mid c\rho \text{ is a controller for } H_A \text{ on } A\}$. We will call any element of T a *controller of H on A* . Again, given an element $c \in H$ which is a controller of H on A , we can write elements of H in a unique c -form. I.e., if $g \in H$, and c is a controller for H on A , then there is an integer k so that $g = c^k \overset{\circ}{g}$, where $\overset{\circ}{g}$ will not realize either end of A .

2.6 Towers

We define and study objects called towers which will give us a geometric criterion on derived length.

Given a group $G \leq PL_o(I)$ and a set T of signed orbitals of G , we will say T is a *tower of G* if T satisfies the following properties:

1. T is a chain in the partial order on the signed orbitals of G .
2. For any orbital $A \in O_T$, T has exactly one element of the form (A, g) .

We note that every group $G \in PL_o(I)$ has the empty tower as one of its towers. We will also sometimes create a tower from a given chain in the orbitals of the elements of G , where we will implicitly use a choice function from the set of orbitals of elements of G to G , so that each orbital maps to some element of G which has that orbital. If a non-empty tower is finite, then it will admit an order preserving bijection from a set $\{1, 2, \dots, n\}$ for some positive integer n . We will therefore refer to a finite tower's " i -th" element, by which we mean the image of the integer $i \in \{1, 2, \dots, n\}$ in the tower, extending this, we will also sometimes refer to the tower's "smallest" element, and its "largest" element.

It is a trivial observation that if T is a tower of G for some subgroup $G \in PL_o(I)$ with element (A, g) , then T has no other element of the form (B, g) .

With the definition of tower in place, we have a natural measure of one form of complexity for a group. Given a group $G \leq PL_o(I)$, we define the *depth of G* to be the supremum of the set of cardinalities of the towers of G . Note that groups have depth while towers have height.

We observe that the depth of a group $G \leq PL_o(I)$ is well defined. The set of all towers of G is a nonempty (it contains the empty tower) subset of the power set of the cartesian product of the set of all orbitals of elements of G with the underlying set of the group G . The set of all cardinalities of the set of all towers of G is therefore a nonempty subset of the ordinals, and this set has a supremum. We observe that the depth of the trivial group is

zero, and that the depth of any non-trivial group is greater than zero. In fact, the depth of all abelian groups is one, as we will see shortly.

The following is a good exercise to help the reader become familiar with these concepts. In exploring this, it is helpful to recall that W_i is a balanced group (it is solvable with derived length i , but F is known to be non-solvable, so W_i cannot contain a copy of F) and to observe that α_1 is actually a consistent controller for W_2 on the orbital $(0, 1)$.

Remark 2.21 *The depth of W_2 is two.*

where here we mean the realization of W_2 given in the introduction.

We extend the notions of depth and height to other objects.

Let T be a tower of $PL_o(I)$. We will call the cardinality of T its *height*, using the simple descriptive *infinite* if T has an infinite cardinality. If there is an order preserving injection from $\mathbf{N}$ to T , then we will say T is *tall*, and if there is an order preserving injection from the negative integers to T , then we will say T is *deep*. If T is both deep and tall then we will say T is a *bi-infinite tower*, and we note that there will be an order preserving injection from the integers to the tower. We will occasionally replace an infinite tower of one of these three types with the image of the implied injection without comment, so that we might refer to a tall, deep, or bi-infinite tower as countable, and refer to the “next” element, etc., when this will not effect the result of an argument.

Given a group $G \leq PL_o(I)$, and an orbital A of an element $g \in G$. We will define the *depth of A in G* to be the supremum of the heights of the finite towers which have their smallest element having the form (A, h) for some element $h \in G$. If the depth of A is an infinite ordinal, we will simply say that A is *deep in G* . Symmetrically, we define the *height of A in G* to be the supremum of the heights of the finite towers which have their largest element having the form (A, h) for some element $h \in G$. If the height of A is an infinite ordinal, we will simply say that A is *high in G* .

The following are immediate from the definitions.

Remark 2.22 1. *Any subset of a tower is a tower.*

2. *If T is a tower for some group $G \leq PL_o(I)$, and $(A, g) \in T$, and if h is an element of G with orbital A , then the set $(T \setminus \{(A, g)\}) \cup \{(A, h)\}$ is also a tower for G .*
3. *If T is a tower for some group $G \leq PL_o(I)$, and $(A, g) \in T$, and if n is a non-zero integer, then $(T \setminus \{(A, g)\}) \cup \{(A, g^n)\}$ is also a tower for G .*
4. *If $g \in G \leq PL_o(I)$, and T is a tower of G , and if $(A, g), (B, g) \in T$, then $A = B$. That is, no signature appears twice in a tower.*
5. *Given a tower T of a group $G \leq PL_o(I)$, the group $H \leq G$ generated by the signatures of T has an orbital A which contains all the orbitals of T .*
6. *Given an element $k \in G \leq PL_o(I)$ and any tower T for G , the set of signed orbitals $T^k = \{(Ak, g^k) | (A, g) \in T\}$ is also a tower for G , where the natural order of the signed orbitals of T^k is equal to the induced order from the signed orbitals of T .*

Given $k \in G \leq PL_o(I)$ and a tower T for G , the tower T^k induced from the tower T by the action of k as discussed in item 6 of Remark 2.22 will be called the *tower conjugate to T by the action of k* . We will also say that the towers are conjugate towers. Conjugacy of towers for G is an equivalence relation on the set of towers for G .

Towers, as defined, are easy to find, but can be difficult to work with. For an arbitrary tower T , there are no guarantees about how other orbitals of signatures of the elements of T cooperate with the orbitals of the tower. We say a tower T is an *exemplary tower* if the following two additional properties hold:

1. Whenever $(A, g), (B, h) \in T$ then $(A, g) \leq (B, h)$ implies the orbitals of g are disjoint from both ends of the orbital B .
2. Whenever $(A, g), (B, h) \in T$ then $(A, g) \leq (B, h)$ implies no orbital of g in B shares an end with B .

The following three lemmas are an indication of the plethora of exemplary towers in $PL_o(I)$, and will be used repeatedly.

Lemma 2.23 *Suppose $H \leq PL_o(I)$, and that $G \leq H$ has imbalanced orbital $A = (a, b)$. Then G admits an exemplary bi-infinite tower E whose orbitals are all in A .*

pf:

A short incorrect proof is that H contains a copy of F and F contains an exemplary bi-infinite tower. The problem is that Theorem 2.9 guarantees a copy of F , but does not guarantee that its generators have one orbital each.

If we find an exemplary bi-infinite tower E' for the projection G_A of G on A , then by replacing the signatures of E' with elements of G which agree with the signatures of E' on A , we can build a new exemplary tower E for G with the same orbitals. Thus, we may assume for the purposes of this argument that G only has orbital A .

Since A is imbalanced for G , there is $g_0 \in G$ so that g_0 has an orbital B_0 which shares an end with A , but g_0 does not realize the other end of A . We will assume that B_0 shares its right end with the right end of A , in particular, $B_0 = (a_0, b)$ for some $a_0 \in A$, and there is $w \in A$ so that if $x \in \text{Supp}(g_0)$ then $w < x$. We will further assume that g_0 moves points to the right on the orbital B_0 , so that all conjugates of g_0 move points to the right on their corresponding orbitals.

By Lemma 2.8 there is an element $\alpha \in G$ so that $r = w\alpha > a_0$. Let $g_k = g_0^{(\alpha^{-k})}$ for all integers k . By construction, given any integer k , the element g_k has rightmost orbital of the form (a_k, b) , where $a_{k+1} < \inf(\text{Supp}(g_k))$.

We will construct a new sequence from the g_k so that the righthand orbitals of the new sequence do not get arbitrarily small, while preserving all of the nesting properties we will need later. Fix $l \in (a_0, r)$.

We now have $a < w < a_0 < l < r < b$. Now for each non-negative integer k , we already have that $a_k < l$, so define $h_k = g_k$ when $k \geq 0$. Since h_0 is already defined, we can define h_{-k} inductively for each $k \in \mathbf{N}$ as the first conjugate of g_{-k} by h_{-k+1}^{-1} which has the property that the rightmost orbital (c_{-k}, b) of h_{-k} contains (l, b) . For each integer k , let c_k represent the left end of the rightmost orbital of h_k . In particular, we have now defined a bi-infinite sequence of functions $(h_k)_{k \in \mathbf{Z}}$ that satisfies the following properties:

1. Given $k \in \mathbf{Z}$, the rightmost orbital of h_k is (c_k, b) .
2. Given $k \in \mathbf{Z}$, $(l, b) \subset (c_k, b)$.
3. Given $k \in \mathbf{Z}$, $c_{k+1} < (\inf(\text{Supp}(h_k)))$.

Note that the support of the element g_{-1} lies to the right of r . For all $k \in \mathbf{Z}$, define $u_k = h_k g_{-1}^{-1}$. Since the h_i are all conjugates of g_0 , and g_{-1} is a conjugate of g_0 , we see that the trailing slopes of all of the h_k are the same. In particular, for each integer k , the fixed set of u_k has a component of the form $[e_k, b)$ where $l < r < e_k < b$. Furthermore, for each integer k , u_k has an orbital C_k of the form (c_k, d_k) where $r < d_k \leq e_k < b$. There may be other orbitals of u_k to the right of d_k and the d_i may not be ordered with respect to i .

Let $v_0 = u_0$, and inductively define, for each positive integer k , an element $v_k = u_k^{h^{m_k}}$ where m_k is the smallest positive integer so that the support of v_{k-1} is fully contained in the orbital $D_k = (c_k, r_k)$ of v_k induced from $C_k = (c_k, d_k)$. Note that such an integer m_k will always exist, since h_k moves points to the right on its rightmost orbital (c_k, b) , and by reference to lemma 2.5.

Now, for each positive integer k , inductively define elements v_{-k} by following the following two step process.

First, find the negative integer n_k of smallest absolute value which has the property that the closure of the support of $v'_{-k} = u_{-k}^{(h^{n_k})}$ is fully contained in the orbital D_{-k+1} of v_{-k+1} induced from $C_{-k+1} = (c_{-k+1}, d_{-k+1})$. At this point v_{-k-1} cannot be created since the orbital of v'_{-k} induced from C_{-k} might not contain the left end of the orbital (c_{-k-1}, b) of h_{-k-1} .

Now replace the elements of the sequences $(u_j)_{j \in \mathbf{Z}}$ and $(h_j)_{j \in \mathbf{Z}}$ which have indices less than or equal to $-k$ by the conjugate of each such element by $h_{-k}^{n_k}$. Note $n_k < 0$. This will do nothing to the element h_{-k} of the sequence $(h_j)_{j \in \mathbf{Z}}$, but all terms of $(h_j)_{j \in \mathbf{Z}}$ with $j < -k$ will have their orbitals extended leftward by $|n_k|$ iterates of h_k^{-1} . Now define $v_{-k} = u_{-k}$. At any inductive stage k , note that by construction, all of the left ends of the orbitals of the h_j , for indices $j < -k$, are inside the orbital D_{-k} , so this inductive definition makes sense. For each integer k , the closure of the support of v_{k-1} is a subset of this orbital D_k . In particular, $E = \{(D_k, v_k) | k \in \mathbf{Z}\}$ is an exemplary bi-infinite tower for G , with all orbitals D_k in A , and where the index of E respects the natural order on the signed orbitals of E . $\diamond$

In the balanced group case, we have another way to sometimes find bi-infinite, exemplary towers:

Lemma 2.24 *Suppose H is a balanced subgroup of $PL_o(I)$ and H has orbital A . If h is an element of H so that h realizes the ends of A inconsistently, then H admits an exemplary bi-infinite tower T with the orbitals of T all contained in A .*

pf:

We will find such a tower for the projection of H on A , then replacing the signatures of our tower with elements of H which agree with our signatures over A will create an exemplary tower for H whose orbitals are all in A . Thus, we shall assume that H has only the orbital A for our discussion below.

Let $h \in H$ so that h realizes the ends of A inconsistently. By replacing h with its inverse, if necessary, we can assume that h moves points to the left on its first orbital and moves

points to the right on its last. Let $F_h = \text{fix}(h) \cap A$ represent the (non-empty) fixed set of h in A .

By lemma 2.20, there is an element g_0 in $\hat{G}$ which has an orbital B_0 which fully contains the set F_h . Let $r = \inf(F_h)$ and $s = \sup(F_h)$, so that $F_h \subset [r, s] \subset B_0 \subset \bar{B}_0 \subset A$. We may assume g_0 moves points to the right on B_0 . There is a smallest positive integer n_1 so that the orbital B_1 of $g_1 = g_0^{(h^{n_1})}$ induced from B_0 by the action of h^{n_1} contains the closure of the support of g_0 , since repeated conjugation of g_0 by h increases the size of the orbital B_0 so that the ends of B_0 approach the ends of A . In particular, we can inductively define a bi-infinite sequence $((B_i, g_i))_{i \in \mathbf{Z}}$ of signed orbitals of H by the property that $g_i = g_{i-1}^{(h^{n_i})}$ where n_i is the smallest positive integer so that the closure of the support of g_{i-1} is fully contained in the orbital B_i induced from the orbital B_{i-1} by the action of h^{n_i} (more formally, for negative integers i , one defines g_i from g_{i+1} by saying that $g_i = g_{i+1}^{(h^{-n_{i+1}})}$ where $-n_{i+1}$ is the largest negative integer so that the closure of the support of g_i is contained in B_{i+1} where this last makes sense since repeated conjugation by h^{-1} moves points in the first and last orbitals of h arbitrarily close to the closed interval $[r, s]$).

$T = \{(B_i, g_i) | i \in \mathbf{Z}\}$ is now an exemplary bi-infinite tower for H whose indexing follows the natural ordering of the signed orbitals of T . $\diamond$

In the following recall that S_T is the set of signatures of a tower T .

Lemma 2.25 *Suppose G is a balanced group and G has a tower T of height n for some positive integer n . If B_n is the orbital of $H_n = \langle S_T \rangle$ which contains the orbitals of T , then G has an exemplary tower E of height n whose orbitals are contained in B_n .*

pf:

Let us write $T = \{(A_i, g_i) | 1 \leq i \leq n, i \in \mathbf{N}\}$ where the indexing of T respects the natural ordering of the signed orbitals of T .

For each positive integer k in $\{1, 2, \dots, n\}$, define the following objects:

$$T_k = \{(A_i, g_i) | 1 \leq i \leq k, i \in \mathbf{N}\}, \\ H_k = \langle S_{T_k} \rangle.$$

Furthermore, for each such k , let B_k be the orbital of H_k which contains the orbitals of the tower T_k for H_k , and note that each end of B_k is realized by some signatures from T_k , since H_k is finitely generated. Further, since G is balanced, each signature of T_k which realizes one end of B_k must also realize the other, and so for each k , some signature of T_k must realize both ends of B_k in B_k . But now, by Lemma 2.15 we know that the projection of the group H_k on the orbital B_k has a controller c_k for the orbital B_k .

There are now two cases to analyze.

First, if c_k is inconsistent, then the signature of T_k which realizes both ends of the orbital B_k must do so inconsistently, and so, by Corollary 2.24, H_k will admit an exemplary bi-infinite tower F whose orbitals all are contained in the orbital $B_k \subset B_n$. F will admit an exemplary subtower E of height n all of whose orbitals will be contained in B_n , and we would be finished.

Second, if c_k is consistent, then the signatures of T_k which realize both ends of B_k do so in a consistent manner. But now, by Lemma 2.16, each signature of T_k that realizes both ends of B_k actually realizes B_k , and at least one signature does so, so that signature must

be g_k , since A_k properly contains the orbitals A_i with $i < k$. In particular, $A_k = B_k$ for each integer $k \in \{1, 2, \dots, n\}$.

Given any orbital B of some signature g_i of T_n , where $B \neq A_i$, we see that there is no index k where B contains an end of any orbital A_k of T . Otherwise, we see that $k > i$, so that the orbital B_k of H_k will be realized inconsistently by some signature g_r with $1 \leq r \leq k$, which we have already assumed does not happen. In a similar fashion, B cannot share an end with any orbital A_k in the orbital A_k . Otherwise, we again have $k > i$. Now g_i and g_k generate a balanced group with orbital C containing A_k and B and having their commonly shared end as an end, and therefore both g_i and g_k share the other end of C as well. At the same time, since g_i cannot realize C ($A_i \cup B \subset C$), we must have that g_i realizes the two ends inconsistently, which we have already ruled out. Therefore, $E = T = T_n$ is already exemplary. $\diamond$

The following indicates that subgroups of $PL_o(I)$ without transition chains of length greater than one are structurally much less complex than general subgroups of $PL_o(I)$.

Lemma 2.26 *If G admits a transition chain of length two, then G admits an exemplary bi-infinite tower.*

pf:

If G is imbalanced, then G admits an exemplary bi-infinite tower by Lemma 2.23, and we are finished, so let us suppose instead that G is balanced.

Suppose G admits a transition chain $\mathcal{C}' = \{(B_1, g_1), (B_2, g_2)\}$. G now must admit a maximal transition chain $\mathcal{C} = \{(A_i, h_i) | 1 \leq i \leq n, i \in \mathbf{N}\}$ using only the signatures g_1 and g_2 , where n is some integer greater than one. Define $H = \langle S_{\mathcal{C}} \rangle$. H is a finitely generated subgroup of G with orbital A containing the orbitals A_i . Now one of g_1 and g_2 must realize the left end of A , and since H is balanced, it must also realize the right end of A . We will assume that g_1 realizes the ends of A for purposes of discussion, since the other case is completely symmetric. But now, since g_1 does not realize A , it must realize both ends of A inconsistently, by Corollary 2.16, and therefore by Lemma 2.24, H (and therefore G) admits an exemplary bi-infinite tower all of whose orbitals are in A . $\diamond$

Corollary 2.27 *If G is a balanced subgroup of $PL_o(I)$ and G admits a tall tower in some orbital A , or G admits a deep tower in some orbital A , or G admits a bi-infinite tower in some orbital A , then G admits an exemplary tall tower in A , or G admits an exemplary deep tower in A , or G admits an exemplary bi-infinite tower in A , respectively.*

pf:

This follows easily from the details of the proof of Lemma 2.25. Suppose T is an infinite non-exemplary tower for a balanced subgroup G , where all the orbitals of T are contained in the orbital A of G . Since T is not exemplary, then we can produce a non-exemplary subtower $P = \{(A_1, g_1), (A_2, g_2)\}$ of T where $A_1 \subset A_2$. Suppose B is the orbital of $G_P = \langle g_1, g_2 \rangle$ that contains A_1 (and therefore A_2 , also note that $B \subset A$). Since P is not exemplary, we must have that some orbital of g_1 contains an end of A_2 or shares an end of A_2 in A_2 . In the first case, G_P admits a bi-infinite exemplary tower in B by Lemma 2.26, which has a subtower of the appropriate type. In the other case, $A_2 = B$, so that A_2 is actually an orbital of G_P .

Now since we are in the second case, we must have that g_1 has some orbital C in A_2 that shares one end of A_2 in A_2 . Since G_P is balanced, we have that g_1 realizes both ends of A_2 from within A_2 . But now g_1 realizes A_2 since G is balanced and the orbital A_2 is realized consistently by g_2 . But this means that $A_2 = A_1$, which contradicts the fact that $A_1 \neq A_2$. $\diamond$

We note in passing that it is an open question as to whether there are any finitely generated, balanced subgroups of $PL_o(I)$ which do not admit transition chains of length two, but which are non-solvable. This can be considered one of the main remaining geometric questions in the theory of solvability of subgroups of $PL_o(I)$. Such a group would be very interesting to examine.

2.7 Derived groups and towers

The following lemma represents the key concept for understanding why solvability and depth are connected. Note that here and in the remainder we use $[a, b] = a^{-1}b^{-1}ab$.

Lemma 2.28 *Suppose G is a subgroup of $PL_o(I)$, n is a positive integer, and A is an orbital of G . If G has a tower of height n whose orbitals are contained in A , then G' has a tower of height $n - 1$ whose orbitals are also contained in A .*

pf: Suppose $T = \{(A_i, g_i) | 1 \leq i \leq n, i \in \mathbf{N}\}$ is a tower of height n all of whose orbitals are contained in the orbital A of G and whose indexing respects the order of the elements of T . By Lemmas 2.23 and 2.25, we can assume that T is an exemplary tower.

For each integer i with $2 \leq i \leq n$ there is a smallest positive integer n_i so that the subset of the support of g_{i-1} which is in A_i is fully contained in a fundamental domain of $g_i^{n_i}$ in A_i , by Lemma 2.5. Define $h_1 = g_1$, and for each integer i with $2 \leq i \leq n$, define $h_i = g_i^{n_i}$. Now define the set $E = \{(A_i, h_i) | 1 \leq i \leq n, i \in \mathbf{N}\}$, which is a exemplary tower. E has a nice property: if i is an integer in $1 \leq i < n$, then the supports of h_i and the supports of $h_i^{h_{i+1}}$ are disjoint in A_{i+1} , so that A_i is an orbital of $[h_i, h_{i+1}]$. For each integer i in $1 \leq i < n$ define $v_i = [h_i, h_{i+1}]$. Noting that $v_i \in G'$, we see that $\{(A_i, v_i) | 1 \leq i \leq n - 1, i \in \mathbf{Z}\}$ is a tower of height $n - 1$ for G' whose orbitals are all in A . $\diamond$

2.8 Geometric classification of solvable groups in $PL_o(I)$

We are now in a position to produce algebraic results by using our geometric tools.

The following is an immediate consequence of Lemma 2.28

Corollary 2.29 *Suppose G is a subgroup of $PL_o(I)$. If G has towers of arbitrary height then G is non-solvable.*

The following is the key lemma towards building a geometric understanding of derived groups in solvable groups of $PL_o(I)$

Lemma 2.30 *If G is a subgroup of $PL_o(I)$ of depth n for some positive integer n , then G' is a subgroup of $PL_o(I)$ of depth $n - 1$.*

pf:

Suppose G is a subgroup of $PL_o(I)$ with depth n . G must be balanced by Lemma 2.23, and G must have no transition chains of length two by Lemma 2.26.

Since $n > 0$, G is not the trivial group, and so G has at least one orbital. Let $A = (a, b)$ be an orbital of G . Note that if

$$\Upsilon = \{B \mid B \text{ is an orbital of some element of } G, B \subset A\},$$

then $A = \cup_{B \in \Upsilon} B$. But G has no transition chains of length two or the depth of G would be infinite, so A can be written as a union of a chain of properly nested orbitals of elements of G . Taking these orbitals, paired with appropriate signatures, we create a tower T whose height is bounded above by n . Let the height of T be m , and let $T = \{(A_i, g_i) \mid 1 \leq i \leq m, i \in \mathbf{Z}\}$. But now, by construction, A is the union of the orbitals A_i , all of which are contained in A_m , so that A_m must actually be A , and no element $g \in G$ has an orbital B properly containing A_m (or $A = A_m$ would not be an orbital of G). Thus, (A_m, g_m) is a signed orbital of depth one, and the orbitals of G are precisely the orbitals of depth one for G . (Note that since G admits no transition chains, any signed orbital of depth one for G is automatically an orbital of G .)

Let $g, h \in G$, and consider the element $[g, h] \in G'$. Let Γ be the set of all orbitals of g and h . Suppose $[g, h]$ has m orbitals, where m is some positive integer, and let $\{A_i \mid 1 \leq i \leq m, i \in \mathbf{N}\}$ be the orbitals of $[g, h]$ in left to right order. Both g and h fix a , so the slope of the leftmost affine component of $[g, h]$ that intersects A is the product of the slopes of the leftmost affine components of g^{-1} , h^{-1} , g , and h in A , which product is one. In particular, $[g, h]$ cannot realize A , so no orbital of $[g, h]$ is an orbital of G , and so no tower for G' contains an orbital of depth one for G , and thusly, all towers of G' can have height at most $n - 1$.

By the last paragraph, we see that the depth of G' is at most $n - 1$. By Lemma 2.28 G' has a tower of depth $n - 1$, so the depth of G' is actually $n - 1$. $\diamond$

We are now ready to prove our main geometric result, that given a non-negative integer n , a subgroup H in $PL_o(I)$ is solvable with derived length n if and only if H has depth n .

Proof of Theorem 1.9:

If the derived length of a group is n , then it must contain a tower of height at least n , otherwise by Lemma 2.30 and the fact that a depth zero group is trivial, the derived series will terminate too soon. But if G has a tower of height greater than n , then again by Lemma 2.30, the n -th derived group $G^{(n)}$ of G will admit a tower of height at least one, and so $G^{(n)}$ will not be trivial. $\diamond$

The next lemma is a technical lemma that we will use in completing our proof of Theorem 1.1.

Lemma 2.31 *If G is a solvable subgroup of $PL_o(I)$ with derived length n , generated by a collection Γ of elements of $PL_o(I)$ which each admit exactly one orbital, and where no generator can be conjugated by an element of G to share an orbital with a different generator, then G is isomorphic to a group in the class $\{1\}\mathcal{P}$ with derived length n .*

pf: Before getting into the main body of the proof, note that the hypotheses imply that each generator is the only generator with that orbital, and that no element orbital is a union of element orbitals that do not realize the original orbital.

We now enter the main body of the proof. We will proceed by induction on n .

If $n = 0$ then G is the trivial group, and $G \in \{1\}\mathcal{P}$. If $n = 1$ then G is abelian, and in particular, there can be at most countably many generators in Γ , all of which have disjoint support, so that G is isomorphic with a countable (or finite) direct sum of $\mathbf{Z}$ factors.

Now let us suppose that $n > 1$ and that the statement of the lemma is correct for any such solvable group with derived length $n - 1$. Let X represent the generators in Γ whose orbitals are all depth 2 or deeper. Let Y be the set of elements in Γ whose orbitals have depth 1. We note in passing that the cardinality of Y is at most countably infinite, and that the collection of orbitals of the elements of Y actually form the orbitals of the group G . We will assume that all of the elements in Y move points to the right on their orbitals. We can partition the elements in X into sets P_y , where the y index runs over the elements in Y , and where an element of X is in P_y if and only if that element's orbital is contained inside the orbital of y . Given $y \in Y$, if P_y is empty, then define $H_y = \langle y \rangle$. Otherwise, let $\gamma \in P_y$, and suppose that γ has smallest depth possible for the elements in P_y , and that γ has orbital $A = (a, b)$. y has a fundamental domain $D_y = [a, ay)$, and each element of P_y may be conjugated by some power of y so that the resultant element's orbital lies in the fundamental domain D_y (if some element, β , conjugates to contain a in its orbital, then either that conjugate has that its orbital fully contains the orbital A , which is impossible by our choice of γ as having a minimal depth orbital of the orbitals of all the elements in P_y , or the signed orbitals of γ and of the conjugate of β form a transition chain of length two, which is impossible since G is solvable). We can now replace P_y by the conjugates of the original P_y found above, and the group generated by the new P_y with y will be identical to the group generated by the old P_y with y . However, now that all of the elements of P_y have supports in the same fundamental domain of y , we have that $H_y = \langle P_y, y \rangle$ is isomorphic to $K_y \wr \mathbf{Z}$, where $K_y = \langle P_y \rangle$. But K_y is a solvable group of precisely the type mentioned in the hypotheses of the lemma, with derived length k less than n , so that K_y is isomorphic to a group in $\{1\}\mathcal{P}$ with derived length $k < n$, which implies that H_y is isomorphic to a group in $\{1\}\mathcal{P}$ (being the result of a group in $\{1\}\mathcal{P}$ being wreathed with a $\mathbf{Z}$ factor on the right) with derived length $k + 1 \leq n$. But this argument holds for every y in Y so that

$$G \cong \bigoplus_{y \in Y} H_y$$

and since all of the groups in this countable direct sum have derived length less than or equal to n (and at least one of them has derived length n), we see that G is isomorphic to a group in $\{1\}\mathcal{P}$ with derived length n . $\diamond$

The following is commonly used without comment in the remainder, since we often work in the situation where a group in $PL_o(I)$ is balanced and admits no transition chains of length two.

Remark 2.32 *Suppose $G \leq PL_o(I)$ and G is balanced and does not admit transition chains of length two. If (A, g) and (B, h) are signed orbitals of G and $A \cap B \neq \emptyset$ then either $\overline{A} \subset B$, $\overline{B} \subset A$, or $A = B$.*

pf:

Suppose (A, g) and (B, h) are signed orbitals of G and $A \cap B \neq \emptyset$. Suppose $A = (a, b)$ and $B = (c, d)$. Let $H = \langle g, h \rangle$, and let $C = (e, f)$ be the orbital of H which contains $A \cup B$. We will assume without loss of generality that $a \leq c$. Since a is not interior to any orbital of h , we must have that $e = a$. If $b < d$ then in order for $\{(A, g), (B, h)\}$ not to form a transition chain of length two, we must have that $c = a = e$. In this case $f = d$, since d cannot be in any orbital of g without creating a different transition chain of length two. Now h will realize the orbital C of H consistently, and g realizes the left end of C in C , so g must then also realize the right end of C . This means that g realizes both ends of C in C and must be consistent on the ends of C since h is consistent. By Lemma 2.16 we see that g cannot have any fixed points in C . But then $b = d$, which contradicts our assumption that $b < d$. In particular, we must have that $d \leq b$. If $d = b$ then by reasoning similar to the previous case, we must have that $a = c$, so that $A = B$. In particular let us assume that $d < b$. Again, if $a = c$ we will have that A is an imbalanced orbital of H , so we must have that $c > a$. In particular, we have shown that if $a \leq c$ then either $\overline{B} \subset A$ or $A = B$. By a symmetric argument, if $c \leq a$ then either $\overline{A} \subset B$ or $A = B$. In particular, either $A = B$, $\overline{A} \subset B$, or $\overline{B} \subset A$. $\diamond$

One great tool for technical analysis of a subgroup G of $PL_o(I)$ is the *split group* of G . It is motivated by the hypotheses of Lemma 2.31. Suppose G is a subgroup of $PL_o(I)$, and let Γ be the maximal set of elements of $PL_o(I)$ which all have single orbitals, where if $\gamma \in \Gamma$, then γ is identical to an element h of G on γ 's orbital. The group $\langle \Gamma \rangle$ is the split group of G . Note that $G \leq \langle \Gamma \rangle$.

Here is another technical lemma, which we use for our main result below.

Technical Lemma 2.33 *Suppose G is a balanced subgroup of $PL_o(I)$ that admits no transition chains of length two, that has derived length t for some $t \in \mathbf{N}$, and that has split group H . If (A, h) is a signed orbital of H , then there is $g \in G$ so that (A, g) is a signed orbital of G .*

pf:

Let us denote by Γ the maximal set of elements of $PL_o(I)$ which all have one orbital, so that if $\gamma \in \Gamma$ then γ is identical to an element of G over γ 's orbital, so that $H = \langle \Gamma \rangle$.

For each non-negative $k \in \mathbf{Z}$, let $P(k)$ be the statement that if $\gamma = \gamma_1 \gamma_2 \cdots \gamma_k$ is a product of elements of Γ of length k for some non-negative $k \in \mathbf{Z}$, and if A is an orbital of γ , then there is an element α of Γ which also has orbital A .

We will prove that $P(k)$ is true for all non-negative $k \in \mathbf{Z}$ via induction, at which point we will have proven the lemma.

If $k = 0$ the statement $P(0)$ is vacuously true. If $k = 1$ the statement $P(1)$ is trivially true. The key statement of the induction proof occurs when $k = 2$. Suppose $\gamma = \gamma_1 \gamma_2$, where $\gamma_1, \gamma_2 \in \Gamma$, and let A be an orbital of γ . We have two possible cases, which will be exhaustive as a consequence of Remark 2.32.

1. γ_1 and γ_2 share an orbital C with $\overline{A} \subset C$, or
2. at least one of γ_1 and γ_2 has the orbital A , while the other has orbital B with $B \subset A$ or $B \cap A = \emptyset$.

Let us suppose it is the first case. Then there are elements f and g in G with f identical to γ_1 on C , and g identical to γ_2 on C . The product fg is an element in G with orbital A so that fg is identical to γ on A , so in this case, $P(2)$ is true.

The second case trivially satisfies the statement $P(2)$, so we have shown $P(2)$ is true in all cases.

Therefore let us assume that $k > 2$ and that we know that $P(m)$ is true for all integers m with $0 \leq m < k$.

Let $\gamma = \gamma_1\gamma_2 \cdots \gamma_m$ be a product of m elements from Γ with an orbital A . Let $g_{m-1} = \gamma_1\gamma_2 \cdots \gamma_{m-1}$. The product $g_{m-1}\gamma_m$ has orbital A , so either both g_{m-1} and γ_m share an orbital C with $A \subset C$, or at least one of g_{m-1} and γ_m has orbital A . If either of these two elements has orbital A , we are done, since we have found a shorter product of elements of Γ that produces the orbital A (possibly length one). So let us assume that instead that $\bar{A} \subset C$, and C is the orbital of both g_{m-1} and γ_m . In this case, there is a smallest index j with $1 \leq j \leq m$ so that γ_j has orbital D with $A \subset D$ and all the orbitals of the γ_i are contained in or disjoint from D . We note that we can permute the order of the products, moving γ_j all the way to the left and replacing the γ_i that γ_j moves past with a conjugate element, which we can still find in Γ (since the orbital of γ_j always contains or is disjoint from the orbitals of the elements we are passing across, the new conjugate elements are guaranteed to be in Γ , which is not the case if the orbital of γ_j was interior to the orbital we were trying to pass across), so that the total number of elements in the product stays constant, and the resulting product still is the same element of H . Repeat this procedure with the smallest index in the set of indices $\{2, 3, \dots, m\}$ so that we place this new element in the second slot of the product that produces γ . γ_1 and γ_2 now have the largest orbitals of the orbitals of the γ_i which contain A . It is immediate therefore that D is also the orbital of γ_2 , else γ will have orbital D which contains $\bar{A}$. There are elements g_1 and g_2 in G which agree with γ_1 and γ_2 over D , so that the element g_1g_2 in G has every orbital of $\gamma_1\gamma_2$ as an orbital of g_1g_2 , so that these orbitals are all realized by elements in Γ . Now, if $\gamma_1\gamma_2$ has orbital D , then by our induction hypothesis, we are finished, since g_1g_2 would be an element of G which behaves as the product $\gamma_1\gamma_2$ over D , so that we could write γ as a shorter product of elements in Γ . Therefore let us assume that the product $\gamma_1\gamma_2$ does not have D as an orbital. In this case all of the orbitals of $\gamma_1\gamma_2$ are interior orbitals of D . Replace the product $\gamma_1\gamma_2$ by the product $\tau_1\tau_2 \cdots \tau_r$ where the τ_i are elements in Γ that agree with $\gamma_1\gamma_2$ on the r element orbitals of $\gamma_1\gamma_2$. Since these orbitals are disjoint, the ordering in this product is immaterial. We can even remove any τ_i with an orbital that does not contain the boundary of A , since such a τ_i will not effect whether A is a resultant orbital of the overall product. In particular, we will retain at most two τ_i , so that the total length of the product did not change. However, there are fewer elements in the product with orbital D . In particular, we can now repeat this process. At each stage, the number of elements in the product that realize the largest orbital containing A , or the size of that largest orbital, is decreasing, while the number of elements in the product does not go up. Since G is solvable, this process must eventually halt, since the largest element orbital of the new γ_i cannot keep shrinking using orbitals in Γ , or G would possess an infinite tower. $\diamond$

Corollary 2.34 *Suppose G is a subgroup of $PL_o(I)$ and let H be the split group of G , then the derived length of G equals the derived length of H .*

pf:

If G is solvable, then G must be balanced and admits no transition chains of length two, so the Lemma 2.33 applies, and therefore given any tower T of H we can find a tower with the same orbitals in G . The other direction is immediate.

If G is non-solvable, then both G and H have towers of arbitrary height. $\diamond$

The following lemma, and its corollary, complete our proof of both Lemma 2.3 and Theorem 1.1. Note that the corollary is simply a restatement of Lemma 1.4.

Lemma 2.35 *If G is a solvable subgroup of $PL_o(I)$ with derived length n , then G is isomorphic to a subgroup of a group H in $\{1\}\mathcal{P}$ so that H has derived length n .*

pf:

Suppose that $n \in \mathbf{Z}$ with $n \geq 0$ and that G is a solvable subgroup of $PL_o(I)$ with derived length n . We see immediately that G is balanced and admits no transition chains of length 2.

Let H be the split group of G , and let Γ be the collection of one-bump generators of H , that is, Γ is the largest collection of one-bump elements in $PL_o(I)$ which satisfies that if $g_A \in \Gamma$ with orbital A , then there is a $g \in G$ with orbital A , and the element g_A equals g on A . We know from lemma 2.34 that H is also solvable with derived length n .

Let X_1 represent the set of signed orbitals of H with depth 1. For each orbital A in O_{X_1} , there is a non-empty set of controllers of H for A in the set Γ . Let $\phi_1 : O_{X_1} \rightarrow \Gamma$ represent a function that associates to each orbital in O_{X_1} a controller of H for that orbital which moves points to the right on the orbital. Let $Y_1 = O_{X_1}\phi_1$ be the image of ϕ_1 . We note that each pair of elements in Y_1 have disjoint support, and trivially, that no element of Y_1 can be conjugated by an element of H to share an orbital with a different element of Y_1 . Now, by the definition of controller, Y_1 consists of a set of generators in Γ sufficient so that the set $\Gamma_1 = Y_1 \cup (\Gamma \setminus S_{X_1})$ generates H .

For each $y \in Y_1$, let A_1^y represent the orbital of y . We may partition the elements of Γ_1 into sets P_1^y indexed by the set Y_1 so that $\gamma \in P_1^y$ if the orbital of γ is contained in A_1^y . Now let X_2^y represent the set of signed orbitals of elements in P_1^y with depth two in H . If X_2^y is not empty, let (A_2^y, γ_2^y) be an element in X_2^y , and let a_y be the left end of the orbital A_2^y . Since the orbitals of elements of H are always orbitals of G by Lemma 2.33, we see that H admits no transition chains of length two. In particular, we can use y to conjugate every signed orbital of X_2^y into the fundamental domain $[a_y, a_y y)$ to produce the set D_2^y , since all of these orbitals can be conjugated to fit in the fundamental domain (else we can get an orbital to have a_y in its interior, creating a transition chain of length two). And likewise we can conjugate every signed orbital in P_1^y of depth greater than two into the fundamental domain as well, producing the set T_2^y . The collections of conjugates in the fundamental domain have nice properties:

1. If two elements of D_2^y have orbitals that non-trivially intersect each other, then they actually have identical orbitals.
2. $H_{A_1^y} = \langle y, S_{D_2^y}, S_{T_2^y} \rangle$.

Let $\phi_2^y : O_{D_2^y} \rightarrow S_{D_2^y}$ be a function that picks for each orbital of depth two in $O_{D_2^y}$ a controller that moves points to the right for that orbital as before. Let Y_2 be the union of all the images of the functions ϕ_2^y across the Y index set, so that we have now picked an controller that moves points to the right on its orbital for every conjugacy class of depth two orbitals of H (note that a conjugate of a controller is also a controller), so that if $\Gamma_2 = (\Gamma_1 \setminus (\cup_{y \in Y} S_{X_2^y})) \cup Y_2$ then $H = \langle \Gamma_2 \rangle$.

In a like fashion we can inductively proceed to pick sets of controllers, one for each conjugacy class of element orbital of depth i , where i is an index less than or equal to n , in exactly the same fashion as discussed for forming the set Y_2 above. This process will steadily improve the sets of generators Γ_i , so that finally Γ_n will be a set of generators for H where each generator in Γ_n has exactly one orbital, and where no generator can be conjugated in G to share an orbital with another generator in Γ_n . In particular, by Lemma 2.31, H is isomorphic to a group in $\{1\}\mathcal{P}$ with derived length n , and therefore G is isomorphic to a subgroup of a group in the class $\{1\}\mathcal{P}$ with derived length n . $\diamond$

Corollary 2.36 *If G is solvable in $PL_o(I)$ of derived length n , then G embeds in G_n .*

pf:

Suppose G is a subgroup of $PL_o(I)$ with derived length n . Lemma 2.35 guarantees that G embeds in a group $H \in \{1\}\mathcal{P}$ with derived length n . But now, Lemma 2.2 guarantees that H embeds in G_n , so that G must embed in G_n . $\diamond$

3 Algebraic classification of non-solvable subgroups of $PL_o(I)$

Our primary result is Theorem 1.2, which states that W is a subgroup of any non-solvable subgroup of $PL_o(I)$. This result is easier than what we actually prove in this section. See the next subsection for a list of the main results.

It is immediate from Theorem 1.9 that a subgroup H of $PL_o(I)$ is non-solvable if and only if H admits towers of arbitrary height. If H admits towers of arbitrary height then we know from section 2 that H admits exemplary towers of arbitrary height. If we consider the realization of the groups W_i in the introduction, then we might suspect that the signatures of an exemplary tower of height n will generate a group isomorphic to W_n , and that the signatures of a countably infinite exemplary tower will generate one of the three groups $(\imath\mathbf{Z})^\infty$, $(\imath\mathbf{Z})^\infty$, and $(\mathbf{Z})^\infty$. This is false, as individual signatures of an exemplary tower may have multiple orbitals; we are only guaranteed that the ends of the orbitals of the exemplary tower are arranged nicely with respect to the set of all orbitals of the signatures of the tower. Nonetheless, this suspicion has still some kernal of truth in it, and so it allows us a toe-hold on what work needs to be done (cleaning up the non-tower orbitals) in order to find towers so that their signatures generate groups that we can analyze.

Suppose that H is a non-solvable subgroup of $PL_o(I)$, then we can informally outline our approach in this section in terms of H as follows.

1. Show that given an exemplary infinite tower for H , we can pass to an even better infinite tower so that the group generated by the signatures of the new tower is one we can recognize. This takes two generic steps.
 - (a) Clean the tower further so that the orbitals of any signature in the tower which are contained in an orbital of the tower are arranged nicely in that orbital.
 - (b) Clean the tower further still so that the orbitals of the signatures which are contained away from the orbital of H which supports the tower are arranged nicely with respect to each other.
2. Show that even in circumstances where we only know that we can find exemplary towers of arbitrary height in H , with reasonable extra information we can conclude there are actually exemplary infinite towers in H .

3. Show that W embeds in any of the groups generated by the process outlined above when H admits an infinite exemplary tower.
4. Show that if H does not admit infinite towers, but is non-solvable, then W embeds in H as a subgroup.

In order to carry out the two types (a) and (b) of tower cleaning mentioned above, we will need to carry out a technical analysis describing the results of building two types of commutators, so recall that in these investigations we use the commutator symbol as follows:

$$[a, b] = a^{-1}b^{-1}ab = a^{-1}a^b.$$

The main points of the outline above will be visible in the statements of results below, although one should not be misled into believing that the statements below precisely follow the development of the proof of the primary result.

3.1 Statement of non-solvability results

First we find conditions under which we can guarantee that we can find a subgroup isomorphic to one of the wreath products mentioned in the introduction.

Theorem 3.1 *Any subgroup H of $PL_o(I)$ which admits a tall tower contains a subgroup isomorphic to $(\mathbf{Z}\wr)^{\infty}$.*

Theorem 3.2 *Any subgroup H of $PL_o(I)$ which admits a deep tower contains a subgroup isomorphic to $(\wr \mathbf{Z})^{\infty}$.*

Corollary 3.3 *If H is a subgroup of $PL_o(I)$ and H admits transition chains of length two then H has subgroups isomorphic to both $(\wr \mathbf{Z})^{\infty}$ and $(\mathbf{Z}\wr)^{\infty}$.*

We also can find one of these wreath products in any finitely generated non-solvable subgroup of $PL_o(I)$.

Theorem 3.4 *Any finitely generated subgroup of $PL_o(I)$ with towers of arbitrary height contains one of $(\mathbf{Z}\wr)^{\infty}$ or $(\wr \mathbf{Z})^{\infty}$ as a subgroup.*

However, there are non-solvable subgroups of $PL_o(I)$ which do not contain any of the three main infinite wreath products mentioned in the introduction.

Lemma 3.5 *Neither $(\wr \mathbf{Z})^{\infty}$, nor $(\mathbf{Z}\wr)^{\infty}$, nor $(\wr \mathbf{Z}\wr)^{\infty}$ embed in W .*

Although converse statements are false, as stated below.

Lemma 3.6 *W embeds in both $(\wr \mathbf{Z})^{\infty}$ and $(\mathbf{Z}\wr)^{\infty}$.*

Therefore, the following lemma completes a proof of Theorem 1.2.

Lemma 3.7 *If H is a non-solvable subgroup of $PL_o(I)$ which does not admit infinite towers then H contains a subgroup isomorphic to W .*

and from this we have an immediate consequence, based on the easy fact that W admits no finite index solvable subgroups.

Corollary 3.8 *Virtually solvable subgroups of $PL_o(I)$ are solvable.*

3.2 Improving exemplary towers

In this section we build technical results guaranteeing that we can often find towers whose signatures generate groups that we can analyze.

The first result is a simple refinement of Lemma 2.20 using Lemma 2.8.

Technical Lemma 3.9 *Suppose H is a balanced subgroup of $PL_o(I)$ and $H = \langle \alpha, \beta \rangle$ for some two elements $\alpha, \beta \in PL_o(I)$. Suppose further that A is an inconsistent orbital of H and α realizes both ends of A while β realizes neither. There is a conjugate γ of β in H which has an orbital $B \subset A$ so that the fixed set of α in A is contained in B .*

Pf: Let F_α represent the fixed set of α in A , and let $x = \inf(F_\alpha)$ and $y = \sup(F_\alpha)$. By Lemma 2.8, since A is an orbital of H , there is $\theta \in H$ so that $x\theta > y$. By the continuity of θ , there is $x_1 < x$ so that $x_1 > y$ as well. Let $z = x_1\theta^{-1}$ so we have

$$z < z\theta = x_1 < x < y < x_1\theta < x\theta$$

Now since F_α is in the orbitals of β , we see that β has an orbital $C = (r, s)$ so that $r < x < s$. There is a power $k \in \mathbf{Z}$ so that $r\alpha^k = q < z$. Now, $\beta_1 = \beta^{\alpha^k}$ has orbital $D = (q, t)$ induced from C by the action of α^k , and D satisfies $q < z < x < t$. Set $\gamma = \beta_1^\theta$. γ has orbital $B = (u, v)$ induced from D by the action of θ on β_1 , and $u = q\theta < x < y < t\theta = v$.

◇

The following lemma is more involved, and plays a key role in the proof of the lemma following immediately after.

Technical Lemma 3.10 *Suppose H is a balanced subgroup of $PL_o(I)$ and $H = \langle \alpha, \beta \rangle$ for some two elements $\alpha, \beta \in PL_o(I)$. If H has an inconsistent orbital A , and β realizes the ends of no orbitals of H , then there are elements α_1 and β_1 in H so that if $H_1 = \langle \alpha_1, \beta_1 \rangle$ then H_1 , α_1 , and β_1 satisfy the following properties:*

1. A is an inconsistent orbital of H_1 .
2. β_1 realizes no ends of any orbital of H_1 .
3. Every inconsistent orbital of H_1 is the union of the orbitals of a transition chain of length three whose first and last orbitals are orbitals of α_1 , and whose second orbital is an orbital of β_1 .
4. α_1 moves points to the left on its leading orbital in each of the inconsistent orbitals of H_1 .

pf:

Set α_1 to be either α or α^{-1} , so that α_1 moves points to the left on its leading orbital B contained in A .

Suppose $n \in \mathbf{N}$ and H has n inconsistent orbitals. There are non-negative integers $r, s \in \mathbf{Z}$ so that $n = r + s$, with $r > 0$, where r is the number of the inconsistent orbitals of H that have α_1 moving points to the left on its leading interior orbitals in these inconsistent

orbitals. Let $\mathcal{B} = \{B_i \mid 1 \leq i \leq r\}$ represent the collection of inconsistent orbitals of H where α_1 moves points to the left on its leading orbital in each of these orbitals, indexed from left to right. Let $\mathcal{C} = \{C_j \mid 1 \leq j \leq s\}$ represent the other inconsistent orbitals of H , and note that if $s = 0$, then this could be an empty collection.

By Technical Lemma 3.9, for each orbital B_i in $\mathcal{B}$ there is an element γ_i in H , which is a conjugate of β , so that the fixed set of α_1 in B_i is contained in a single orbital of γ_i . Likewise, for each orbital C_j in $\mathcal{C}$ there is an element θ_j in H , which is a conjugate of β , so that the fixed set of α_1 in C_j is contained in a single orbital of θ_j .

Firstly, inductively replace each element γ_i , for $i > 1$, by a conjugate of γ_i by a high negative power of α_1 so that whenever $k, i \in \mathbf{N}$, with $1 \leq i < k \leq r$, we have that the closure of the union of all of the orbitals of γ_k that intersect the orbitals of γ_i nontrivially in B_i is actually fully contained in the single orbital of γ_i that contains the fixed set of α_1 in B_i . Note that conjugating by high negative powers of α_1 , we are pushing the orbitals of the conjugates closer to the exterior boundaries of the fixed set of α_1 in each B_i , so that the last sentence is possible.

For each $i \in \{1, 2, \dots, r\}$, γ_i has an orbital D_i that contains the fixed set of α_1 in B_i , as well as the closure of all of the orbitals of γ_k in B_i for all $i < k \leq r$.

We will inductively define a sequence of elements $(\rho_i)_{i=1}^r$. Then, modulo replacing some of the γ_i by their conjugates by more negative powers of α_1 , the ρ_i will have the following properties:

1. $\rho_1 = \gamma_1$.
2. For all indices $i > 1$, ρ_i will be either a conjugate of ρ_{i-1} by some power of γ_i , or $\rho_i = \rho_{i-1}\gamma_i$.
3. For all indices i , ρ_i will have an orbital E_i in B_i that fully contains the fixed set of α_1 in B_i .
4. If $i < r$, the orbital E_i of ρ_i will contain the closures of the orbitals of γ_j in B_i for all integers j with $i < j \leq r$.
5. If $i > 1$, for each integer j with $1 \leq j < i$, ρ_i will have E_j as one of its orbitals.

Firstly, set $\rho_1 = \gamma_1$, and $E_1 = D_1$. By construction, ρ_1 satisfies the five inductive properties. If $r = 1$, we are done. If not, suppose that k is an integer so that $1 < k \leq r$ and for all $i \in \{1, 2, \dots, k-1\}$ we have that ρ_i is defined and satisfies the five defining properties of the induction. Our analysis now breaks into two cases.

If ρ_{k-1} has an orbital F_k containing either end of D_k , then there is some integer j so that $\rho_k = \rho_{k-1}^{\gamma_k^j}$ will have orbital E_k induced from F_k by the action of γ_k^j so that E_k will contain the fixed set of α_1 in B_k , as well as the closure of all of the orbitals of γ_j in B_k for integers j with $i < j \leq r$ (if $k < r$).

If ρ_{k-1} does not have an orbital F_k containing either end of D_k , then we have to handle the case where ρ_{k-1} has orbitals in D_k that share ends with D_k separately before continuing.

If ρ_{k-1} has orbitals in D_k that share ends with D_k then replace γ_k and all later γ_j with conjugates of these elements by a high negative power of α_1 so that D_k either has an end

contained in an orbital of ρ_{k-1} , or shares no end with an orbital of ρ_{k-1} , and repeat the whole inductive definition of ρ_k .

If ρ_k is still undefined, then set $\rho_k = \rho_{k-1}\gamma_k$. Note that since ρ_{k-1} has no orbitals in D_k that share ends with D_k , the product $\rho_k = \rho_{k-1}\gamma_k$ realizes both ends of D_k consistently, and therefore realizes D_k consistently since H is balanced. Therefore define $E_k = D_k$ and note that ρ_k actually has E_k as an orbital.

At this point, ρ_k and E_k are both defined, and we can continue with our main argument. Note that E_k contains the closure of all of the orbitals of all of the γ_i for $i > k$, and that for each integer j in $1 \leq j \leq k-1$, the closure of the orbitals of γ_k in B_j are fully contained in the orbital E_j , so that ρ_k will have E_j as an orbital as well. Now by construction, ρ_k satisfies the five defining properties of the induction.

We now examine the element ρ_r . Observe that the element ρ_r contains an orbital E_k in each B_k where the fixed set of α_1 in B_k is fully contained in E_k . ρ_r is constructed as a sequence of products using various γ_i 's and conjugates of γ_i 's so ρ_r realizes no end of any orbital of H , but is an element of H .

In an entirely analogous fashion, if $s > 0$, then we can find one element ψ_s in H which realizes no end of any orbital of H and which contains an orbital F_i in each $C_i \in \mathcal{C}$ which contains the fixed set of α_1 in that C_i .

There is a positive integer p so that $\rho = \rho_r^{\alpha_1^p}$ has the properties that follow:

1. For each integer $i \in \{1, 2, \dots, r\}$, the closure of the orbitals of ψ_s in B_i is actually contained in the orbital G_i of ρ induced from E_i by the action of α_1^p .
2. For each integer $i \in \{1, 2, \dots, s\}$, the closure of the orbitals of ρ in C_i is actually contained in the orbital F_i of ψ_s .

This follows since for each orbital B_i of $\mathcal{B}$, the lead orbital of α_1 in B_i has the property that α_1 moves points to the left there (and therefore moves points to the right on the trailing orbital of α_1 in B_i), and for each orbital C_i in $\mathcal{C}$, the lead orbital of α_1 in C_i has the property that α_1 is moving points to the right there (and therefore α_1 moves points to the left on its trailing orbital in C_i).

We note in passing that the orbitals G_i of ρ contain the orbitals E_i of ρ_r , and therefore the fixed set of α_1 in the B_i .

Now there is a power q of ψ_s so that the element $\beta_1 = \rho^{\psi_s^q}$ will have the following nice properties:

1. For each integer $i \in \{1, 2, \dots, s\}$, the orbitals of β_1 in C_i have trivial intersection with the fixed set of α_1 in C_i .
2. For each integer $i \in \{1, 2, \dots, r\}$, β_1 will have the orbital G_i which contains the fixed set of α_1 in B_i .

The first property follows since the orbitals of ρ in the C_i are contained in the orbitals F_i of ψ_s , and so the conjugation of ρ by a high power of ψ_s will throw these orbitals off of the fixed set of α_1 in the C_i . The second property follows since the orbitals of ψ_s are fully

contained in the orbitals G_i of ρ in the B_i , so that conjugation of ρ by ψ_s to any power will not change these orbitals.

We observe that by generating the group $H_1 = \langle \alpha_1, \beta_1 \rangle$, we succeed in producing the promised group, since $A = B_k$ of some integer $k \in \{1, 2, \dots, r\}$ is an inconsistent orbital of H_1 , β_1 realizes no ends of any orbital of H_1 , and on each inconsistent orbital D of H_1 , D is the union of a transition chain of length three of the form $\{(L, \alpha_1), (G_k, \beta_1), (R, \alpha_1)\}$ where L is an orbital of α_1 where α_1 moves points to the left, and L contains the left end of G_k , and R is an orbital of α_1 where α_1 moves points to the right, and R contains the right end of G_k , and where G_k is one of the orbitals G_i of the definition of β_1 .

◇

We already know that imbalanced groups contain copies of Thompson's group F , which itself contains many copies of $(\wr \mathbf{Z})^\infty$, (see [2] for details of this last point), so in our investigations, we will generally work under the assumption that the groups we are examining are balanced.

The next result is an example of what we are aiming for when we "improve" exemplary towers, and depends on the previous technical lemma. We will use this result in the proofs of Lemma's 3.16 and 3.17 to immediately reduce to the special case of examining exemplary towers whose signatures generate a group which admits no transition chains of length two.

Lemma 3.11 *If G is balanced and G contains transition chains of length two, then there is an exemplary bi-infinite tower $E = \{(B_i, \beta_i) \mid i \in \mathbf{Z}\}$ for G where the group $H = \langle S_E \rangle$ admits no transition chains of length two.*

pf:

Since G admits transition chains of length two, we can find two elements α and β which form the set of signatures of a transition chain of length two. Let $K = \langle \alpha, \beta \rangle$.

The orbitals of K are the components of the union of the orbitals of α and β . Some of these orbitals may be consistent orbitals for K , so that at least one of α or β realize these orbitals. The other orbitals are inconsistent, and are formed by the union of a subcollection of orbitals of α and orbitals of β . Each such subcollection admits a transition chain of length two. A chief feature of the inconsistent orbitals is that one of α or β must realize both ends of any particular such orbital, since K is balanced. Since K admits transition chains of length two, at least one of the orbitals of K is inconsistent.

We are going to analyze the orbitals of K still further. Any particular orbital of K has one of six types, the first three are consistent, and the last three are inconsistent:

1. (Type AB) Both α and β consistently realize this orbital.
2. (Type Ab) α consistently realizes this orbital, but not β .
3. (Type aB) β consistently realizes this orbital, but not α .
4. (Type ab) Both α and β inconsistently realize both ends of this orbital.
5. (Type ab) α inconsistently realizes both ends of this orbital, but β realizes neither end of this orbital.

6. (Type $a\bar{b}$) β inconsistently realizes both ends of this orbital, but α realizes neither end of this orbital.

We know that K has at least one orbital, let us call it A , of type $\underline{ab}$, $\underline{ab}$, or $a\bar{b}$, and we will assume without meaningful loss of generality that A has one of the first two types. Let F_a represent the union of the fixed sets of α that are contained in the orbitals of K of type $\underline{ab}$, type $\underline{ab}$, and $a\bar{b}$. F_a is non-empty, and is entirely contained in the orbitals of β . In particular, by Remark 2.5 there is a $N_1 \in \mathbf{N}$ so that for all $k \in \mathbf{N}$ with $k \geq N_1$ we have $F_a\beta^k \cap F_a = \emptyset$ in orbitals of type $\underline{ab}$ and $\underline{ab}$ (in orbitals of type $a\bar{b}$, the interior components of F_a are moved off of themselves). Similarly, let S represent the support of α in the orbitals of K of type aB , then there is $N_2 \in \mathbf{N}$ so that for all $k \geq N_2$, we have $S\beta^k \cap S = \emptyset$. Let $N = \max(N_1, N_2)$. Considering the other direction, let F_b represent the fixed set of β in the orbitals of K of type $a\bar{b}$. Since F_b is contained in the support of α by definition, there is $M \in \mathbf{N}$ so that for all $j \geq M$ we have that $F_b\alpha^j \cap F_b = \emptyset$. Now let $j \geq M$, and let $k \geq N$, and define $\beta_1 = [\alpha^j, \beta^k]$. We observe that the fixed set of α in the orbitals of K of type $\underline{ab}$ and $\underline{ab}$ is contained in the orbitals of β_1 . Note also that the components of F_a in the orbitals of K of type $a\bar{b}$ which do not realize any end of an orbital of K are all contained in the support of β_1 . The fixed set of β contained in the orbitals of K of type $a\bar{b}$ is also contained in the support of β_1 , since any such point is moved off of F_b by α^{-j} , then moved by β^{-k} , then moved to someplace different (from its start) by α^j , and finally, β^k cannot move the resultant point to its original location in the fixed set of β . Now observe that the orbitals of β_1 are either disjoint from S , or else are components of S where α^j behaves as the inverse of β_1 .

We now consider the group $K_1 = \langle \alpha, \beta_1 \rangle$, and we will consider the orbitals of K_1 under the same classification as the orbitals of K , where we replace β by β_1 in that classification.

It is immediate to see that K_1 still has all the orbitals of K of type $\underline{ab}$, and that the type of these orbitals is unchanged. It is also immediate by construction that the orbitals of K of type $\underline{ab}$ are also orbitals of K_1 , although they are now of type $\underline{ab}$. The orbitals of K of type Ab are also orbitals of K_1 of type Ab , but the orbitals of K of type aB are now replaced by a collection of interior orbitals (all lying properly in the union of the orbitals of K of type aB), each of which is an orbital of type aB that is actually disjoint from the support of α , or else of type AB , where α^j and β_1 behave as inverses on these orbitals. The orbitals of K of type AB are now of type Ab , and may have trivial intersection with the support of β_1 (if, in fact, α and β commuted on these orbitals).

If B_1 is an orbital of K of type $a\bar{b}$, then B_1 is not an orbital of K_1 . In this case K_1 admits a new collection of orbitals properly contained in B_1 .

We first consider the case where β is moving points to the right on its leading orbital in B_1 (and therefore is moving points to the left on its trailing such orbital). We will suppose k was chosen large enough so that the closure of the union of the orbitals of $\beta^{-k}\alpha^j\beta^k$ that are contained in orbitals of β in B_1 is actually contained in the orbitals of α (and therefore of α^j) which contain components of the fixed set of β . Note that any interior orbital of β in B_1 is contained in the union of the orbitals of α^j and $\beta^{-k}\alpha^j\beta^k$. Therefore, there are three possible varieties of resulting orbitals of K_1 in B_1 : firstly, of type AB , where β_1 actually behaves as α^{-j} on these orbitals (there may be several of these), secondly, of type AB , where there is only one such orbital, and it contains the fixed set of β , or thirdly, of type $\underline{ab}$, where there

is one of these if the previous variety did not occur, and it contains the fixed set of β in this case. We will assume k was chosen large enough so that these properties of transformation are preserved over all orbitals of K of type $a\underline{b}$ where β is moving points to the right on its leading relevant orbitals.

In the case of the orbitals of K of type $a\underline{b}$ where β moves points to the left on its leading relevant orbitals. The results depend heavily on the nature of α in these individual orbitals. To clarify the discussion, let us suppose that B is such an orbital, and discuss the possibilities that arise from the behaviour of α and β on B .

Firstly, let us suppose that α has an orbital that contains the fixed set of β in B . In this case, let us suppose k and j were chosen large enough so that the entire support of α is contained inside a single fundamental domain of the single orbital of $\beta^{-k}\alpha^j\beta^k$ that contains the fixed set of β in B . In this case, the group K_1 possibly has several orbitals in B , all of type aB . One of these orbitals contains all of the support of α in B , and all of the rest are orbitals of β_1 which contain no orbitals of α and are therefore of type aB with trivial intersection with orbitals of α .

Now let us suppose that α has more than one orbital in B that contains a component of the fixed set of β . The first and last such orbitals of α in B must have that α behaves inconsistently on these orbitals, otherwise it is easy to create an imbalanced subgroup of K_1 . So now there are two further cases.

Let us suppose that α moves points to the right on its first orbital in B which contains a component of the fixed set of β , and therefore moves points to the left on the last such. In this case K_1 has only one orbital in the domain B , call it C , which is again of type $a\underline{b}$. The closure of C is contained in B , and β_1 moves points to the right on its leading orbital in C and moves points to the left on its trailing orbital there.

Now let us suppose α moves points to the left on its leading orbital in B that contains a component of the fixed set of β , and therefore moves points to the right on its trailing orbital in B which contains a component of the fixed set of β , the group K_1 again has some pure orbitals (type aB) plus precisely one orbital C in B , which is again of type $a\underline{b}$, and this time, β_1 will move points to the left on its leading orbital in C and will move points to the right on its trailing orbital in C .

The result of all of this analysis is the following, we can choose j and k so that the group K_1 has orbitals of the following types:

1. AB

Note that in this case α and β_1 commute on this orbital, except in the case possibly generated from orbitals of type $a\underline{b}$ where b moves points right on its leading orbital.

2. Ab

3. aB

Note here that the behavior of α on this orbital is as the identity, unless this orbital is contained in an orbital of K of type $a\underline{b}$, in which case α may have non-trivial support in this orbital.

4. $\underline{ab}$

Note that orbitals of this type are always contained in orbitals of K of type $a\bar{b}$ where β moves points to the right on its first relevant orbital.

5. $\underline{ab}$

Since these are the certain result of an orbital of type $\underline{ab}$ or of an orbital of type $\underline{ab}$ of K , we see that K_1 will have at least one of these.

6. $a\bar{b}$

These orbitals all have the property that whenever β_1 moves points to the left on its leading orbital in these orbitals, then α moves points to the left on its leading orbital of the orbitals that contain a component of the fixed set of β_1 .

In particular, we can repeat this process to create a new element β_2 using α and β_1 , and therefore a new group $K_2 = \langle \alpha, \beta_2 \rangle$. K_2 improves on K_1 since all of its orbitals of type $a\bar{b}$ have both β_2 and α moving points to the left on their important leading orbitals. In particular, K_2 may still have orbitals of type $\underline{ab}$, and of type AB (although here α and β_2 will commute on these orbitals). K_2 may have orbitals of type Ab , but its orbitals of type aB will all have the property that α is the identity over these orbitals, while K_2 will certainly have orbitals of type $\underline{ab}$. Repeating the process one more time to create an element β_3 and a subgroup $K_3 = \langle \alpha, \beta_3 \rangle$ produces a group whose orbitals are much easier to describe. K_3 will have no orbitals of type AB since K_2 had no orbitals of type aB or $a\bar{b}$ that could produce these orbitals (the types exist, but not with the right subflavors of α and β_2 to generate these offspring). K_3 may have orbitals of type Ab , but it will have no orbitals of type aB , since the orbitals of type aB in K_2 had α behaving as the identity there, and K_2 had no orbitals of type $a\bar{b}$ with β_2 moving points to the left on its first sub-orbital D while α was moving points to the right on its orbital containing the right end of D . K_3 will have no orbitals of type $\underline{ab}$, since K_2 had no orbitals of type $a\bar{b}$ with β_2 moving points to the right on its first orbital in the orbitals of K_2 of this type. K_3 will have at least one orbital of type $\underline{ab}$, and may have several orbitals of the type $a\bar{b}$, but all of these last will have β_3 moving points to the left on its leftmost orbitals in these orbitals, and α will also move points to the left on its first orbitals containing the right ends of β_3 's leftmost orbitals in these orbitals of type $a\bar{b}$ of K_3 .

Now, the orbitals of K_3 are well understood, and the behaviors of β_3 and α on these orbitals are also well understood. We now consider the subgroup K_4 generated by α and $\beta_4 = [\alpha^{-j}, \beta_3^k]$, where j and k are chosen as in the previous process (note the negative index on α). The point of this is that now the orbitals of K_4 will admit no orbital of type $a\bar{b}$ with β_4 moving points to the left on its first orbital. Now replacing K_4 with $K_5 = \langle \alpha, \beta_5 \rangle$ where $\beta_5 = [\alpha^j, \beta_3^k]$ where j and k are chosen as before produces a group with no orbitals of type $a\bar{b}$, repeating one more time to generate β_6 and K_6 in the same fashion that we generated K_1 from K produces a group whose orbitals are only of types Ab and $\underline{ab}$.

Let us consider the orbital A of K . A is also an orbital of K_6 , and it is of type $\underline{ab}$. We will now replace K by K_6 and β by β_6 so that K has an orbital of type $\underline{ab}$ and all of its orbitals are of type $\underline{ab}$ and Ab .

Suppose K has n orbitals of type $\underline{ab}$, and let $\mathcal{O} = \{A_i \mid 1 \leq i \leq n, i \in \mathbf{N}\}$ represent this collection, where the indices respect the left to right order of the orbitals. By construction

we know that $n \geq 1$. Apply Technical Lemma 3.10 (above) to replace α and β by new elements, and replace K by the new group generated by the new α and β so that β still realizes no end of any orbital of K , and A_1 is still an orbital of type $\underline{ab}$, but where every maximal transition chain (of length greater than one) formable using α and β has length three (naturally α provides the leading and trailing orbitals for any such chain), and where α moves points to the left on all of its leading orbitals in orbitals of type $\underline{ab}$ for K (and therefore moving points to the right on its trailing such intervals).

Define $\gamma_0 = \beta$. For each $i \in \mathbf{N}$, inductively define $\gamma_i = \gamma_{i-1}^{\alpha^{k_i}}$ where k_i is chosen large enough so that in each orbital of K of type $\underline{ab}$, the closure of all of the orbitals of γ_{i-1} is contained in the single orbital of γ_i that contains the fixed set of α in that particular orbital of K of type $\underline{ab}$, and so that in any particular orbital X of K of type Ab , the set of orbitals of γ_i lie between an end of X (already chosen by the direction α moves points) and the orbitals of γ_j in X for all j with $0 \leq j < i$. Note that this k_i exists, since α moves points to the left on all of its leading orbitals in orbitals of K of type $\underline{ab}$.

Similarly, define the γ_i for negative integers i by setting $\gamma_i = \gamma_{i+1}^{\alpha^{k_i}}$ where k_i is chosen large enough (in the negative direction) so that in any orbital of K of type $\underline{ab}$ the closure of all of the orbitals of γ_i is contained in the orbital of γ_{i+1} that contains the fixed set of α in that orbital of K , and so that the orbitals of γ_i in orbitals of K of type Ab are disjoint from all of the orbitals of the γ_k for $k > i$. Note that this last is possible in these orbitals of K of type Ab , despite the fact that there are infinitely many such orbitals to avoid, since we are now conjugating by α to negative powers, so that orbitals of the γ_i for negative i all lie on the other side of the orbitals of γ_0 compared to the orbitals of γ_i with $i > 0$.

By construction, if i and j are integers, and $i < j$, then any orbital C of γ_i that intersects an orbital D of γ_j has the property that $\overline{C} < D$. Furthermore, $\overline{D}$ is then contained in an orbital of γ_k for every integer $k > j$. In particular, for each integer i there is $m_i \in \mathbf{N}$ so that whenever B is an orbital of γ_i , all of the orbitals of γ_{i-1} in B are actually contained in a single fundamental domain f_{m_i} of $\gamma_i^{m_i}$ in B . Inductively, we see that all of the orbitals of γ_k in B for any integer $k < i$ are actually also contained in f_{m_i} . Define $\beta_i = \gamma_i^{m_i}$.

The group $H = \langle \{\beta_i \mid i \in \mathbf{Z}\} \rangle$ now admits no transition chains of length two. For every integer i , let B_i represent the orbital of β_i that contains the fixed set of α in A_1 . By construction, the B_i all exist, and are nested (so that $B_i \subset B_j$ whenever $i < j$) so that $E = \{(B_i, \beta_i) \mid i \in \mathbf{Z}\}$ is an exemplary bi-infinite tower for H whose indexing respects the ordering on the signed orbitals of E . $\diamond$

Suppose $h, k \in PL_o(I)$ and they satisfy the properties that whenever A is an orbital of h and B is an orbital of k , and $A \cap B \neq \emptyset$, then either $A = B$, or $\bar{A} \subset B$, or $\bar{B} \subset A$. (Note that if H is a balanced subgroup of $PL_o(I)$ which admits no transition chains, then any two elements of H will satisfy these conditions.)

Under these conditions, we will say that h and k satisfy the *mutual efficiency condition* if given any orbital of C of h that properly contains an orbital of k , then the support of k in C is contained in a single fundamental domain of h in C , and the symmetric condition that whenever D is an orbital of k that properly contains an orbital of h , then the support of h in D is contained in a single fundametal domain of k in D .

Note that the intitial containment conditions on the orbitals of h and k above occur for any two elements in any balanced subgroup of $PL_o(I)$ with no transition chains of length two.

The following remark follows easily from Lemma 2.5 and the nature of the orbital alignments of the elements of the hypotheses, and is left to the reader.

Remark 3.12 *If h and k are elements of a balanced subgroup of $PL_o(I)$ that admits no transition chains of length two, then there are positive integers m and n so that h^m and k^n satisfy the mutual efficiency condition.*

We will use this fact heavily in the remainder.

The following remark is a simple exercises in the calculus that partially determines the orbitals of a product of two elements of a balanced subgroup of $PL_o(I)$ without transition chains of length two.

Remark 3.13 *Suppose h and k are elements of a subgroup H of $PL_o(I)$, where H is balanced and admits no transition chains of length two.*

1. *Suppose h has an orbital A and k has an orbital B , and $A \subsetneq B$, then hk has orbital B .*
2. *Suppose h has an orbital A and k has an orbital B , and $B \subsetneq A$, then hk has orbital A .*
3. *If C is an orbital of hk , then either C is an orbital of h , and k does not have an orbital properly containing C , or C is an orbital of k and h does not have an orbital properly containing C , or both h and k have orbital D which contains C .*

pf:

To see the first point, since H admits no transition chains of length two, no end of B can be contained in an orbital of h , so B is an orbital of $\langle h, k \rangle$. But k realizes B consistently, so that any element of $\langle h, k \rangle$ which realizes one end of B must realize B . Therefore since A is properly contained in B , h must not realize any end of B . But therefore the product hk is non-trivial near the ends of B in B , so hk must actually realize B .

The argument for the second point is similar to the argument for the first point, and will not be given here.

To see the third point, if neither h nor k have an orbital containing C , then some point x in C is not in the support of either h or k (since H admits no transition chains), so that the product hk cannot have orbital C . Therefore one of h and k must have an orbital D containing C . In which case the first two points imply all three cases of the conclusion to the third point.

◇

We will also use the following construction.

Construction 1 (Double Commutator Operation) *Given two elements $h, g \in PL_o(I)$ we can construct a third element $[g, {}_2h] = [[g, h], h]$, which we will refer to as the double commutator of g and h .*

Double commutators are nice to understand in the setting of a balanced subgroup of $PL_o(I)$ with no transition chains of length two.

Lemma 3.14 *Let $g, h \in H$, where H is a balanced group in $PL_o(I)$ with no transition chains of length two. Suppose further that g and h satisfy the mutual efficiency condition. If $f = [g, {}_2h]$, then f has the following properties:*

1. *Every orbital of g properly contained in an orbital of h is an orbital of f .*
2. *Every orbital of f is properly contained in an orbital of h that contains (perhaps not properly) an orbital of g .*

pf:

Suppose A is an orbital of g properly contained in some orbital B of h . The element $[g, h] = g^{-1}h^{-1}gh$ can be thought of as the product of g^{-1} with the conjugate g^h . Since the support of g in the orbital B of h is fully contained in a single fundamental domain of h in B , the orbitals of g^h are completely disjoint from the orbitals of g in B . In particular the product $[g, h]$ has all the orbitals of g , where $[g, h]$ acts as g^{-1} on these orbitals, and a full disjoint copy of these orbitals where the action of $[g, h]$ on these orbitals is as g^h . Now the element $[g, {}_2h] = ([g, h])^{-1}h^{-1}[g, h]h$ can be thought of as the product of $([g, h])^{-1}$ with the conjugate $[g, h]^h$. But the only orbitals of $([g, h])^{-1}$ which intersect the orbitals of $[g, h]^h$ are precisely the orbitals of g^h , which are disjoint from the original orbitals of g . Thus, since $[g, h]^{-1}$ has all of the orbitals of g , $[g, h]^h$ does not have these orbitals, hence the product $[g, {}_2h]$ has the orbital A . In particular, we have proven the first point of the lemma.

Now let A be some orbital of $f = [g, {}_2h]$. A is either disjoint from the orbitals of h , properly contains an orbital of h , or is contained in an orbital of h .

Suppose $A \cap \text{Supp}(h) = \emptyset$. In this case, let us think of f as the product of two elements of H ,

$$f = [[g, h], h] = (h^{-1})^{[g, h]} \cdot h.$$

In the situation that H is balanced and contains no transition chains of length two, any orbital of the product of two elements in H is actually an orbital from one or the other of the two elements of the product, or is properly contained in an orbital B which is common to the two elements of the product, by Remark 3.13. In this case, since A is disjoint from the orbitals of h , A must actually be an orbital of $(h^{-1})^{[g, h]}$. But then $A = B[g, h]$ for some orbital B of h^{-1} . Now A is disjoint from B (B is also an orbital of h), so in particular, B and A are contained in a single orbital E of $(h^{-1})^g h$, which means E is actually an orbital of $(h^{-1})^g$. This implies in turn that $(A \cup B) \subset Dg$, where D is another orbital of h^{-1} . But now B and D are orbitals of h contained in an orbital F of g , and B is in the image of the orbital D of h under the function g . But every orbital of h in F is thrown off the support of h by the action of g by assumption, providing a contradiction, so that our initial assumption that $A \cap \text{supp}(h) = \emptyset$ must be false.

Now suppose A is an orbital of f which properly contains an orbital of h . We see immediately then that A is an orbital of $(h^{-1})^{[g, h]}$. But then A is either an orbital of $[g, h]^{-1}$ or an orbital of $h^{-1}[g, h]$, or is contained properly in a common orbital B of both of these elements. In the first and second cases, A would actually be an orbital of both elements, since h^{-1} does not have any orbitals in A near the ends of A . But the slopes of $[g, h]^{-1}$ and $h^{-1}[g, h]$ near the ends of A in A are inverse, so that A cannot actually be an orbital of the product $[g, h]^{-1} \cdot h^{-1}[g, h]$. In particular, A must be properly contained in some common

orbital B of both $[g, h]^{-1}$ and $h^{-1}[g, h]$. But then B is an orbital of $h^{-1}[g, h]$ induced from an orbital of $(h^{-1})^g$ by the action of h , which cannot move the ends of B (h has an orbital in A , and therefore in B , and so has no orbital containing the ends of B), so that B is actually an orbital of $(h^{-1})^g$. But every orbital $(h^{-1})^g$ is disjoint from the orbitals of h^{-1} , so that B could not contain an orbital of h .

From the above, we now see that A is contained in an orbital B of h , but we do not yet know that B contains an orbital of g . Suppose firstly that B is disjoint from every orbital of g . In this case it is immediate that f is actually trivial on B , so that A cannot be an orbital of f .

Now we know that A is contained in an orbital B of H which non-trivially intersects an orbital C of g . If $\bar{C}$ is contained in B then it is immediate that $\bar{A} \subset B$ and we are done, so suppose instead the either $C = B$ or $\bar{B} \subset C$.

If $C = B$, then again, the slopes of g and h cancel near the ends of B , so that $\bar{A} \subset B$, and again we are done. Therefore let us assume that $\bar{B} \subset C$.

In this case, given any $x \in A \subset B$, we can directly calculate the impact of f on x to see that $xf = x$. Let us write f as a product to see this.

$$\begin{aligned} xf &= x(g^{-1}h^{-1}gh)^{-1}h^{-1}(g^{-1}h^{-1}gh)h = xh^{-1} \cdot (g^{-1}hg) \cdot h^{-1} \cdot (g^{-1}h^{-1}g) \cdot hh = \\ &= x(h^{-1}h^{-1}) \cdot (g^{-1}h^{-1}g) \cdot (hh) = x(h^{-1}h^{-1}) \cdot (hh) = x \end{aligned}$$

In the above the expressions with $\cdot$'s the parentheses have been placed suggestively to help us understand the dynamics of the point x . Any such factor in an above product involving a “ $\cdot$ ” represents a factor with no net effect on the point it acts on.

Thus, if A is an orbital of f , then A is properly contained in an orbital B of h , and B contains an orbital C of g . $\diamond$

3.3 Finding infinite wreath products in groups with infinite towers

Suppose $D = \{(A_i, h_i) \mid i \in \mathbf{N}\}$ is an exemplary tower whose indexing respects the order of the elements so that $H = \langle S_D \rangle$ is a balanced group that admits no transition chains of length two, and so that whenever B is an orbital of h_i for some signature h_i of D , then B is contained in an orbital C of h_{i+1} . We are going to find a subtower of D that satisfies a nice further property.

Suppose B_1 is an orbital of h_1 . Each signature h_i of D has an orbital B_i that contains B_1 . The orbitals B_j are nested as the index increases, but possibly not properly. If there is an $N_1 \in \mathbf{N}$ so that for all $n > N_1$, we have $B_n = B_{n+1}$, then we will call B_1 a *terminal orbital* of D , and (B_1, h_1) a *terminal signed orbital* of D , and we will say that B_1 is *stable after* N_1 . We now extend this language to orbitals of signatures other than h_1 . Given $i \in \mathbf{N}$, call an orbital of h_i *terminal in* D if the orbital is terminal in the subtower of D formed using only the signed orbitals (A_k, h_k) with $k \geq i$. We will call the orbital of any signature of the tower, where the orbital is not a terminal orbital, a *non-terminal orbital*. Observe that non-terminal signed orbitals make good candidates for being bases of new exemplary towers.

We will rely heavily on the following technique in our proof of Lemma 3.16.

Lemma 3.15 (growing subtower) *Suppose $D = \{(A_i, h_i) \mid i \in \mathbf{N}\}$ is an exemplary tower so that $H = \langle S_D \rangle$ is a balanced group that admits no transition chains of length two, and so that whenever B is an orbital of h_i for some signature h_i of D , then B is contained in an orbital C of h_{i+1} . Then we can pass to an infinite subtower E of D so that if J is an orbital of any signature g_i of E , where J is not a terminal orbital of D , then there is an orbital K of g_{i+1} which properly contains J .*

We note by definition that the orbital K will also be a non-terminal orbital of D , and both will be non-terminal in E .

Proof of Lemma:

We now pass repeatedly to infinite subtowers for D , at each stage referring to the new tower that results as D , and re-indexing so that the tower will still have the form $D = \{(A_i, h_i) \mid i \in \mathbf{N}\}$. Let $\mathcal{P} = \{B_i \mid 1 \leq i \leq n_i, i \in \mathbf{N}\}$ represent the n_i orbitals of h_1 that are not terminal, in left to right order. We improve D by passing to a infinite subtower n_i times. Firstly, for B_1 , pass to a subtower of D so that the orbitals of the h_i over B_1 are always properly nested as we progress up the tower. The new tower D still has all the properties that we have listed for the old D , but now the orbitals of the h_i over B_1 actually form a tower over B_1 when we pair them with their signatures. Repeat this process inductively for each of the non-terminal orbitals of D in h_1 . Now we pass to an infinite induction, by repeating the process again, using base signature h_2 , so that we are progressively improving the tower above h_2 so that the non-terminal orbitals of h_2 are each actually the base of an infinite tower using the signatures h_k with $k > 2$ paired with their appropriate orbital containing the relevant orbital of h_2 . We note in passing that the non-terminal orbitals of h_1 are all contained in the non-terminal orbitals of h_2 , and D is already a perfect tower with respect to these operations over the non-terminal orbitals of h_2 which contain non-terminal orbitals of h_1 , so that we will only have to improve D over the non-terminal orbitals of h_2 which do not contain orbitals of h_1 . With these observations in place, we can inductively continue this process at every level of D . We have now defined a new tower D . Given any $i \in \mathbf{N}$, if (A_i, h_i) is a signed orbital of D and B_i is a non-terminal orbital of h_i then for any integer $k > i$ there is an orbital B_k of h_k so that $\bar{B}_i \subset B_k$. Thus, $D_{B_i} = \{(B_k, h_k) \mid k \geq i, k \in \mathbf{N}\}$ is itself an exemplary tower. $\diamond$

The following lemmas are simply restatements (with proofs) of Theorems 3.1 and Theorem 3.2.

Lemma 3.16 *If G is a subgroup of $PL_o(I)$ and G admits a tall tower, then G has a subgroup of the form $(\mathbf{Z})^\infty$.*

pf:

We will assume that G is balanced, as otherwise G contains a subgroup isomorphic to Thompson's group F , which itself has a subgroup isomorphic to $(\mathbf{Z})^\infty$. If G admits transition chains of length two then by Lemma 3.11 G admits an exemplary bi-infinite tower $E = \{(A_i, g_i) \mid i \in \mathbf{Z}\}$ where the indexing respects the order of the signed orbitals, and where the group $H = \langle S_E \rangle$ admits no transition chains of length two. In particular, we can pass to the subgroup K generated by all the signatures with positive index to reproduce the hypotheses of this lemma, with the extra condition that G admits no transition chains of length two, so let us assume that G admits no transition chains of length two.

Let $E = \{(A_i, g_i) \mid i \in \mathbf{N}\}$ be a tall tower for G , where the indexing respects the order on the signed orbitals of E . Since G is balanced, and contains no transition chains of length two, we see that all towers of G are exemplary, and in particular, E is exemplary.

Let $A = \cup_{i \in \mathbf{N}} A_i = (a, b)$. We observe that if B is an orbital of g_i for some i , then B is disjoint from $\{a, b\}$, and that if $B \cap A \neq \emptyset$, then neither a nor b is an end of B . In particular, A is an orbital of $\langle S_E \rangle$.

Now given $\epsilon > 0$ so that $\epsilon < \frac{b-a}{2}$, we see that there is an $N \in \mathbf{N}$ so that for all $n \in \mathbf{N}$ with $n \geq N$, we have that $(a + \epsilon, b - \epsilon) \subset A_n$ since the ends of the A_i must limit to the ends of A . But now, we can construct a monotone strictly increasing, order preserving function, $\phi : \mathbf{N} \rightarrow \mathbf{N}$, so that given any $n \in \mathbf{N}$, all of the orbitals of g_n in A are actually contained in $A_{\phi(n)}$, and since E is exemplary, no orbital of g_n in A actually shares an end with $A_{\phi(n)}$. For any $k \in \mathbf{N}$, let ϕ^k represent the product (via composition) of the function ϕ with itself k times in the monoid of order preserving functions from $\mathbf{N}$ to $\mathbf{N}$. Now define an order preserving function $\theta : \mathbf{N} \rightarrow \mathbf{N}$, defined by the rules that $1 \mapsto 1$ and $n \mapsto \phi^{n-1}(1)$ for each $n \in \mathbf{N} \setminus \{1\}$. Replace E by the exemplary tower formed by the collection $\{(A_{\theta(i)}, g_{\theta(i)}) \mid i \in \mathbf{N}\}$. E now has the property that if $i, k \in \mathbf{N}$ with $i < k$ then all the orbitals of g_i in A are actually in A_k , away from the ends of A_k . For each $n \in \mathbf{N}$, with $n > 1$, let m_n be an integer large enough so that the collection of orbitals of g_{n-1} inside of A_n (which is all the orbitals of g_{n-1} in A) is actually fully contained in a single fundamental domain of $g_n^{m_n}$ in A_n . Define $n_1 = 1$. Improve E by replacing each signature g_n with $g_n^{m_n}$. Now define $H = \langle S_E \rangle$. We note in passing that A is an orbital of H .

We will now improve E further. Define $h_1 = g_1$. Now for each $n \in \mathbf{N}$ with $n > 1$, inductively define h_n via the following four step process.

First, define $k_n = g_n^{r_n}$, where r_n is a positive integer large enough so that whenever B is an orbital of g_n that is also an orbital of h_{n-1} , then the product $h_{n-1}k_n$ still has orbital B .

Second, define $h'_n = h_{n-1}k_n$. Recall from Remark 3.13 that any orbital of h_{n-1} which properly contains an orbital of k_n will now be an orbital of h'_n , and that any orbital of k_n that properly contains an orbital of h_{n-1} will also be an orbital of h'_n .

h'_n now has an orbital containing every orbital of h_{n-1} .

Third, choose positive integer s_n large enough so that every orbital C of $h'^{s_n}_n$ which properly contains orbitals of h_{n-1} actually contains all such orbitals in a single fundamental domain of $h'^{s_n}_n$ on C .

Fourth, define $h_n = h'^{s_n}_n$. The result is that the sequence $(h_i)_{i \in \mathbf{N}}$ of signatures satisfies the following list of properties.

1. For each $n \in \mathbf{N}$, A_n is an orbital of h_n .
2. For each $n \in \mathbf{N}$ with $n > 1$, the orbitals of h_{n-1} in A are all contained inside a single fundamental domain of h_n in A_n .
3. For each $n \in \mathbf{N}$ with $n > 1$, if B is an orbital of h_n which is not disjoint from the orbitals of h_{n-1} , then there are two possibilities.
 - (a) B is also an orbital of h_{n-1} .
 - (b) B properly contains a non-empty collection of orbitals of h_{n-1} in a single fundamental domain of h_n on B .

In particular, we can form the new exemplary tower $D = \{(A_i, h_i) \mid i \in \mathbf{N}\}$.

D still has the properties that $\cup_{i \in \mathbf{N}} A_i = A$, and that A is an orbital of the group $\langle S_D \rangle$. Further, the signatures satisfy all the properties of the last paragraph.

We will now improve D by replacing it with the result of finding a growing subtower, so that any non-terminal orbital of any signature h_i of D is properly contained in a non-terminal orbital of a signature with index one higher.

Our new D is far superior to our old D , but h_1 will still have terminal orbitals, if it had them to begin with. Suppose h_1 does have some terminal orbitals. Then there is $N_1 \in \mathbf{N}$ so that all the terminal orbitals of h_1 are stable for $n \geq N_1$. Compute a new element $k = [h_{N_1+1,2} h_{N_1+2}]$ (note that condition (3) above is equivalent to saying that h_{N_1+1} and h_{N_1+2} satisfy the mutual efficiency condition when all the orbitals of h_j are contained in orbitals of h_{j+1} for any $j \in \mathbf{N}$). k has the following properties.

1. The orbitals of h_{N_1} which contain the terminal orbitals of h_1 are not contained in the orbitals of k .
2. No orbital of h_{N_1} which is also an orbital of h_{N_1+1} is also an orbital of k (these are all terminal orbitals of h_{N_1} since D is the result of using a growing tower operation).
3. All the non-terminal orbitals of h_{N_1} are still properly contained in the orbitals of k since k contains the non-terminal orbitals of h_{N_1+1} .

Now replace k and h_{N_1} by sufficiently high powers of themselves so that they satisfy the mutual efficiency condition and let $h = [h_{N_1,2} k]$. The resulting h has the following properties.

1. h has no orbitals intersecting the terminal orbitals of h_1 .
2. h has all the non-terminal orbitals of h_{N_1}

Now replace h and h_1 by sufficiently high powers of themselves so that they satisfy the mutual efficiency condition, and replace h_1 by $[h_{1,2} h]$. Now replace h_1 and h_{N_1} by sufficiently high powers of themselves so they satisfy the mutual efficiency condition. Build the tower

$$D' = \{(A_1, h_1)\} \cup \{(A_i, h_i) \mid i \geq N_1, i \in \mathbf{N}\}.$$

In this tower, h_1 admits only non-terminal orbitals, every orbital of h_1 is properly contained in a non-terminal orbital of h_{N_1} , and h_1 still has a copy of every non-terminal orbital that it started with. If we re-index the tower D' and call it D again, then it satisfies all the old properties of the tower D found above, but its bottom element (h_1) has nice orbitals. We can now repeat this whole process for the subtower of D starting from level two and up, so that the new h_2 will admit all the non-terminal orbitals that it started with, and other non-terminal orbitals, and also will contain no terminal orbitals. Inductively proceed up the tower D , redefining all of the h_i , so that the new tower D satisfies the following properties.

1. $A = \cup_{n \in \mathbf{N}} A_n$
2. For each $n \in \mathbf{N}$ with $n > 1$, the orbitals of h_{n-1} in A are all contained inside the orbital A_n of h_n .

3. For each $n \in \mathbf{N}$ with $n > 1$, if B is an orbital of h_n which is not disjoint from the orbitals of h_{n-1} , then B contains the closure of the union of the collection of orbitals of h_{n-1} that intersect B .

Now for each index $j \in \mathbf{N}$, inductively replace h_j and h_{j+1} by sufficiently high powers of themselves so that they satisfy the mutual efficiency condition. (This is actually unnecessary by the details of the proof of Lemma 3.14, but it hurts nothing and explicitly guarantees that all adjacent pairs of signatures of the tower D satisfy the mutual efficiency condition.) Note that each signature (except h_1) may be replaced by progressively higher powers of itself twice in this operation, but that once two signatures are mutually efficient, replacing either signature by a higher power of itself will still result in a pair that are mutually efficient.

Now any pair of adjacent signatures of the tower D satisfy the mutual efficiency condition.

For every $n \in \mathbf{N}$, define the subgroup $H_n = \langle h_1, h_2, \dots, h_n \rangle$ of G . Given any two elements $f, g \in H_n$, since the supports of f and g are contained in the support of h_n , and since the support of h_n in any one orbital of h_{n+1} is contained in a single fundamental domain of h_{n+1} in that orbital, we see that $f^{h_{n+1}^j}$ and $g^{h_{n+1}^k}$ have disjoint supports and therefore commute, whenever $j \neq k$. If $j = k$, then the product of the conjugated f and g is equal to the conjugate of the product of f and g . In particular, the group of finite products of conjugates of elements of H_n by h_{n+1} is isomorphic to $\sum_{j \in \mathbf{Z}} H_n$, where the indexing factor j represents the power of h_{n+1} used in the conjugation of the element from H_n under consideration. But we can write any element of H_{n+1} as a product of an integer power of h_{n+1} with a product of conjugates of elements of H_n by integer powers of h_{n+1} ; in short, $H_{n+1} \cong H_n \wr \mathbf{Z}$, where the $\mathbf{Z}$ factor is the subgroup $\langle h_{n+1} \rangle$ of H_{n+1} .

Now, $H_1 \cong \mathbf{Z}$, so $H_2 \cong \mathbf{Z} \wr \mathbf{Z}$, $H_3 \cong (\mathbf{Z} \wr \mathbf{Z}) \wr \mathbf{Z}$, and etc., so that $H_n \cong ((\dots (\mathbf{Z} \wr \mathbf{Z}) \wr \mathbf{Z}) \dots \wr \mathbf{Z}$ where the finite wreath product has n factors of $\mathbf{Z}$. In particular, the ascending union $H = \langle h_1, h_2, \dots \rangle \cong (\mathbf{Z} \wr)^\infty$.

◇

Lemma 3.17 *If G is a subgroup of $PL_o(I)$ and G admits a deep tower, then G has a subgroup of the form $(\wr \mathbf{Z})^\infty$.*

pf:

We will use a similar technique to the proof of Lemma 3.16, although the analysis in this case is much simpler.

We will assume that G is balanced, as otherwise G contains a subgroup isomorphic to Thompson's group F , which itself has a subgroup isomorphic to $(\mathbf{Z} \wr)^\infty$. If G admits a transition chain of length two, then by Lemma 3.11 G admits an exemplary bi-infinite tower $E = \{(A_i, g_i) \mid i \in \mathbf{Z}\}$ where the indexing respects the order of the signed orbitals, and where the group $H = \langle S_E \rangle$ admits no transition chains of length two. In particular, we can pass to the subgroup K generated by all the signatures with negative index to reproduce the hypotheses of this lemma, with the extra condition that G admits no transition chains of length two. In particular, we can assume that G admits no transition chains of length two.

Since G is balanced and admits no transition chains of length two, any tower for G is exemplary. In particular, let $E = \{(A_{-i}, g_{-i}) \mid i \in \mathbf{N}\}$ be an exemplary deep tower for G where the indexing respects the order on the elements of the tower. Improve E by replacing

the signatures of E with sufficiently high powers of themselves so that given any negative integer i , then g_{i-1} and g_i satisfy the mutual efficiency condition.

Let $A = A_{-1} = (a, b)$. Since E is exemplary, we see that A is actually an orbital of the subgroup $H \leq G$, where $H = \langle S_E \rangle$. For all $i \in \mathbf{N}$ with $i > 1$, inductively improve E (induct on increasing $i \in \mathbf{N}$ in the following discussion) by replacing the signatures of E according to the following three step process.

First, let $h_{-i} = [g_{-i,2} g_{-i+1}]$.

Second, define the new g_{-i} to be h_{-i} .

Third, replace the elements g_{-i+1} , g_{-i} , and g_{-i-1} with sufficiently high powers of themselves, so that given any index $j \in \mathbf{N}$, the elements g_{-j} and g_{-j-1} satisfy the mutual efficiency condition (observe that if $i > 3$, then g_{-i+1} and g_{-i+2} will now still satisfy the mutually efficiency condition, since we are only replacing g_{-i+1} by higher powers of itself, and these two signatures were already mutually efficient, a similar argument shows that g_{-i-1} and g_{-i-2} will be mutually efficient after this operation as well).

Since $A_{-i} \subsetneq A_{-i+1}$ for all integers $i > 1$, we see that the resultant set of signed orbitals is still a tower (and with the same order), so that this inductive definition will simply improve our tower E . Observe further that given any $k \in \mathbf{N}$, then the orbitals of g_{-k-1} are all properly contained in the orbitals of g_{-k} .

Define the set $\Gamma_i = \{g_j \mid j \leq i, j \in \mathbf{Z}\}$ for each negative integer i . For each negative integer i , define $H_i = \langle \Gamma_i \rangle$. For such i , the orbitals of H_i are actually the orbitals of h_i , since all orbitals of the elements g_k with $k < i$ are contained in the orbitals of g_i . Furthermore, for any such $i < -1$, the orbitals of g_i are contained in the orbitals of g_{i+1} in such a way that in any individual orbital B of g_{i+1} , the support of g_i in B is actually fully contained inside a single fundamental domain of g_{i+1} on B . In particular, $H_i \cong H_{i-1} \wr \mathbf{Z}$, where the $\mathbf{Z}$ factor comes from the subgroup $\langle g_i \rangle$ of H_i . But now inductively, since each generator generates a group isomorphic to $\mathbf{Z}$, we see that $H_1 \cong (\wr \mathbf{Z})^\infty$.

◇

The following lemma is only surprising in the sense that its proof is somewhat involved. It belongs in this subsection since (as will be shown) finitely generated non-solvable subgroups of $PL_o(I)$ always contain infinite towers. This lemma, together with Lemmas 3.17 and 3.16, completes the proof of Theorem 3.4.

Lemma 3.18 *If H is a finitely generated and H admits towers of arbitrary height, then H admits an infinite tower.*

Pf:

Suppose H is a finitely generated subgroup of $PL_o(I)$ and H admits a towers of arbitrary height. Let n be the smallest integer so that H has a generating set of size n . Let $\Gamma = \{g_1, g_2, \dots, g_n\}$ be a minimal set of generators for H , where the sum of the number of the orbitals of elements of Γ is also minimal amongst all generating collections for H of size n . Since H has towers of arbitrary height, n is at least two, and H has at least one orbital.

If H is imbalanced, then H admits a bi-infinite tower by Lemma 2.23. So we will assume that H is balanced.

If H admits a transition chain of length two then H admits a bi-infinite exemplary tower, so we will assume that H has no transition chains of length two. In particular, if $h, g \in H$,

A is an orbital of g and B is an orbital of h , where $A \cap B \neq \emptyset$, then either $A = B$ or $\overline{A} \subset B$ or $\overline{B} \subset A$ by Remark 2.32.

Now if H has an element with a deep orbital, we can use the orbital nesting properties of Remark 2.32 to inductively build either a tall tower or a deep tower. In particular, we will now assume that no orbital of any element of H has infinite depth. In particular, the depth of each orbital of each element of H is well defined and is finite.

We note that since H admits towers of arbitrary length, it must have elements of arbitrary finite depth.

Given any tower T , let us call the union of its orbitals the support of the tower. In our case, since there are no transition chains of length two, and since H is balanced, the group generated by the signatures of T will have an orbital equal to the support of T . If the support of a tower T is contained in an orbital A of an element of $h \in H$, let us say that A supports T and also that h supports T . If we assume that all towers of H are finite, then they are always supported by their highest signature, and their highest orbital.

We will now inductively define a sequence of finitely generated groups $(H_i)_{i=1}^\infty$ so that $H_1 = H$ and for any $i \in \mathbf{N}$ we have the following properties:

1. $H_{i+1} \leq H_i$.
2. There is $m_i \in \mathbf{N}$ and H_i has a finite generating set $\Gamma_i = \{g_{ik} | k \in \mathbf{N}, 1 \leq k \leq m_i\}$.
3. H_i has an orbital A_i and only the generator g_{i1} in Γ_i realizes A_i .
4. H_i admits a collection X_i of towers of arbitrary height for H_i , where all the towers of X_i contain the signed orbital (A_i, g_{i1}) as their largest element.
5. Every tower of X_k is supported by A_i , whenever $k \in \mathbf{N}$ and $k > i$.
6. The orbital A_{i+1} of H_{i+1} satisfies $\overline{A_{i+1}} \subset A_i$.

First, let $H_1 = H$. Since H_1 is finitely generated, it has only finitely many orbitals. Since every tower for H_1 is supported by one of the orbitals of H_1 , one of the orbitals of H_1 supports towers for H_1 of arbitrary height. Define A_1 to be such an orbital of H_1 . Now since H_1 is balanced, H_1 admits a controller for H_1 on A_1 . Let g_{11} be such a controller for H_1 on A_1 . For each $k \in \mathbf{N}$, where $1 \leq k \leq n$, the generator g_k in Γ has a g_{11} -form $g_{11}^{n_{1k}} h_{1k}$ for A_1 , where the orbitals of h_{1k} in A_1 are all properly contained in A_1 . For each $k \in \{1, 2, \dots, n\}$, define $g_{1(k+1)} = h_{1k}$. Now if we set $m_1 = n + 1$ and define the set $\Gamma_1 = \{g_{1k} | 1 \leq k \leq m_1, k \in \mathbf{N}\}$, we see that the set Γ_1 generates H_1 , but only $g_{11} \in \Gamma_1$ actually realizes the orbital A_1 .

Given any tower of H_1 supported by A_1 , either the tower has a signed orbital of the form (A_1, f) as a top element (there can be none higher) for some element $f \in H_1$, in which case we can replace this element by (A_1, g_{11}) , or the tower does not have such an element, in which case we can just add the element (A_1, g_{11}) to the tower. The resultant tower from either of the two processes above will have height unchanged, or one greater than before, and it will have (A_1, g_{11}) as its top element. In particular, we can do this for each of the towers for H_1 supported by A_1 (which set of towers admits towers of arbitrary height). Now define X_1 to be the set of all towers for H_1 which have top element (A_1, g_{11}) . By our discussion,

we see that the collection X_1 of towers of H_1 with top element (A_1, g_{11}) is a collection with towers of arbitrary height.

We see that H_1 satisfies all of the requirements.

Suppose $(B, \tilde{f})$ is a signed orbital of H_1 of depth two. Since $\tilde{f} \in H_1$, there is an element $f \in H_1$ so that f agrees with $\tilde{f}$ on B , where f admits a shortest length decomposition as a product of elements of Γ_1 and their inverses, out of all elements in H_1 which agree with $\tilde{f}$ on B . Let m be the length of such a shortest length product, and let $f = \prod_{i=1}^m k_i$ be such a product. Now B is also an orbital of f and $f|_B = \tilde{f}|_B$, so we will transfer consideration to the signed orbital (B, f) .

Consider the sequence of functions $(f_j)_{j=1}^m$ defined by the rule

$$\begin{aligned} f_j &= (k_1 k_2 \cdots k_{j-1})^{-1} f (k_1 k_2 \cdots k_{j-1}) = \\ &= (k_1 k_2 \cdots k_{j-1})^{-1} (k_1 k_2 \cdots k_m) (k_1 k_2 \cdots k_{j-1}) = \\ &= k_j k_{j+1} \cdots k_m k_1 k_2 \cdots k_{j-1}, \end{aligned}$$

and the corresponding sequence of signed orbitals $((B_j, f_j))_{j=1}^m$ where for each j the orbital B_j is the signed orbital of f_j induced from B by the action of $\prod_{i=1}^{j-1} k_i$ on f . Note that the depth of each orbital B_j is two, and $B_j \subset A_1$.

For each $j \in \{1, 2, \dots, m\}$, we must have that B_j has nontrivial intersection with an orbital of k_j , or the product decomposition could be shortened to produce a different element g which is identical to f (and therefore $\tilde{f}$) on B .

Since f does not have orbital A_1 , we see that the number of times that g_{11} and the number of times that g_{11}^{-1} appear in the product decomposition of f above must be the same, and since f is not trivial, some other elements from Γ_1 (or their inverses) must appear in the product decomposition of f . Let $S = \{i \in \mathbf{N} | 1 \leq i \leq m, k_i \neq g_{11}^{\pm 1}\}$. By the previous comment, S is not empty, let t represent the size of S . Let $\alpha : \{1, 2, \dots, t\} \rightarrow S$ be a monic increasing function, so that $\alpha(1)$ is the smallest element in S , and $\alpha(t)$ is the largest, ie., α is an ordered indexing of the elements in S .

Let $j \in \{2, 3, \dots, t\}$. By the definition of an induced orbital, we see that $B_j = B k_1 k_2 \cdots k_{j-1}$, and that $B_1 = B$. In particular, for $i \in \{2, 3, \dots, t\}$,

$$B_{(\alpha(i))} = B_{(\alpha(i-1))} k_{(\alpha(i-1))} k_{(\alpha(i-1)+1)} k_{(\alpha(i-1)+2)} \cdots k_{(\alpha(i)-1)},$$

and $B_{\alpha(1)} = B k_1 k_2 \cdots k_{(\alpha(1)-1)}$ if $\alpha(1) \neq 1$ and $B_{\alpha(1)} = B$ if $\alpha(1) = 1$.

Claim: For some $i \in S$, we have that $B_{\alpha(i)}$ is actually an orbital of $k_{\alpha(i)}$, so that (B, f) is conjugate to (B_j, f_j) for some index j , where B_j is actually an orbital of one of the generators of Γ_1 .

Consider $f_{\alpha(1)} = k_{\alpha(1)} \cdot v_1$, where v_1 is the product of the remaining terms of the product decomposition of $f_{\alpha(1)}$ after the first term. Now, $k_{\alpha(1)} \neq g_{11}^{\pm 1}$ by construction. By Lemma 2.32 and Remark 3.13 we must have that $B_{\alpha(1)}$ is either an orbital of $k_{\alpha(1)}$, in which case we have our claim, or else it simply properly contains a smaller orbital of $k_{\alpha(1)}$ and is in fact an orbital of v_1 . (Note, $k_{\alpha(1)}$ cannot have an orbital which contains $B_{\alpha(1)}$ since $B_{\alpha(1)}$ is a depth two orbital contained only in the depth one orbital A_1 .) Suppose therefore that $B_{\alpha(1)}$ is an

orbital of v_1 and that $B_{\alpha(1)}$ properly contains an orbital of $k_{\alpha(1)}$. Both of the products

$$\begin{aligned}\tilde{\beta}_1 &= \prod_{j=\alpha(1)}^{\alpha(2)-1} k_j \\ \text{and} \\ \beta_1 &= \prod_{j=\alpha(1)+1}^{\alpha(2)-1} k_j\end{aligned}$$

will take $B_{\alpha(1)}$ to $B_{\alpha(2)}$, so the conjugate of v_1 by β_1 will produce a new function

$$w_2 = k_{\alpha(2)} \cdots k_m k_1 \cdots k_{(\alpha(1)-1)} k_{(\alpha(1)+1)} k_{(\alpha(1)+2)} \cdots k_{(\alpha(2)-1)}$$

which is like $f_{\alpha(2)}$ in that it still has orbital $B_{\alpha(2)}$, and it is the same product as $f_{\alpha(2)}$ excepting the term $k_{\alpha(1)}$ is deleted from the product. Now we can write $w_2 = k_{\alpha(2)} \cdot v_2$, where v_2 is the product in the definition of w_2 , excepting the first term. As before, $B_{\alpha(2)}$ is either an orbital of $k_{\alpha(2)}$ or it is an orbital of v_2 . If it is an orbital of $k_{\alpha(2)}$ we are finished with the claim, otherwise we will inductively follow this procedure, defining elements w_j and v_j along the way for each $j \in \{2, 3, \dots, t\}$, and checking whether $B_{\alpha(j)}$ is an orbital of $k_{\alpha(j)}$ for each $j \in \{1, 2, \dots, t\}$. If this condition is never satisfied, then $B_{\alpha(t)}$ must be an orbital of v_t . But v_t is a product purely of elements which are all g_{11} and its inverse, where the number of each type is the same, so that v_t is actually the identity! In particular, we must have that for some $i \in S$, B_i is an orbital of k_i , so that (B, f) is conjugate to (B_i, f_i) , where $f_i = k_i \cdot u_i$, where $k_i \in \Gamma_1$, or $k_i^{-1} \in \Gamma_1$, and B_i is an orbital of depth two for H_1 which is also an orbital of k_i . The claim is proven.

With this last claim in hand, we see that the elements of Γ_1 actually contain orbitals of depth two, and there are finitely many of them. Further, each orbital of depth two in A_1 for H_1 is conjugate to some element of this finite list. In particular, every tower for H_1 supported by A_1 can be conjugated to a tower supported by a signed orbital (B, g) of depth two where $g \in \Gamma_1$. Now there are towers of arbitrary height for H_1 , but there are only finitely many signed orbitals of the form (A, g) where $g \in \Gamma_1$, so at least one element of Γ_1 has an orbital of depth two that supports towers of arbitrary height beneath it. Let (A_2, g) be a signed orbital of H_1 , where $g \in \Gamma_1$, and A_2 is an orbital of depth two which supports towers of arbitrary height for H_1 beneath it.

Define

$$\Lambda_2 = \{s \in H_1 \mid s = g_{11}^{-k} h g_{11}^k, \text{ Supp}(s) \cap A_2 \neq \emptyset, h \in \Gamma_1 \setminus \{g_{11}\}, k \in \mathbf{Z}\}.$$

the set of all conjugates of elements of Γ_1 (aside from g_{11}) which have resulting support intersecting nontrivially with A_2 . This set is finite, since after multiple conjugates of any element of Γ_1 (other than g_{11}) by g_{11} , the result has all orbitals near an end of A_1 , away from A_2 .

We will define H_2 to be the group generated by Λ_2 . Note that A_2 is an orbital of H_2 and is realized by a generator in Λ_2 .

Claim:

The collection $\hat{X}_2$ of towers for H_2 all of which have largest element of the form (A_2, h) for various $h \in H_2$ contains towers of arbitrary height for H_2 .

In order to establish this, we show that if (A, f) is a signed orbital of H_1 , where $A \subset A_2$, then there is an element $g \in H_2$ so that A is an orbital of g .

Suppose therefore that (A, f) is a signed orbital of H_1 where $A \subset A_2$. Let $A_f = \{g \in H_1 \mid g|_A = f|_A\}$. Let $g \in A_f$ be an element which can be written as a product of elements in Γ_1 (and their inverses) which uses a minimum number of elements distinct from g_{11} or g_{11}^{-1} in the product. Let $g = \prod_{i=1}^s k_i$ be such a product decomposition, and let t be the number of elements distinct from g_{11} and g_{11}^{-1} in the product (note that $t > 0$). Let $P = |\{i \in \{1, 2, \dots, s\} \mid k_i = g_{11}\}|$ and let $N = |\{i \in \{1, 2, \dots, s\} \mid k_i = g_{11}^{-1}\}|$. Since g must have orbital A , and $\bar{A} \subset A_1$, we see that $P = N$. In particular, by inserting an equal number of elements g_{11} and g_{11}^{-1} in appropriate places, we can think of g as actually being a product of conjugates of elements of $\Gamma_1 \setminus \{g_{11}\}$ (and their inverses) by elements of the form g_{11}^j , where $j \in \mathbf{Z}$. Let $g = \prod_{i=1}^t v_i$ be such an expression for g , formed as described.

If $t = 1$, then we note that $g = v_1$, and that v_1 is a conjugate of an element, h , of $\Gamma_1 \setminus \{g_{11}\}$ (or an inverse of one such), which conjugate has orbital $A \subset A_2$. But now A_2 is an orbital of depth two, and A is of depth at least two, so that $v_1 \in \Lambda_2$ or $v_1^{-1} \in \Lambda_2$. In this case, we see that $g \in H_2$.

Now let us assume that $t > 1$. Let $w = \prod_{i=2}^t v_i$. Now $g = v_1 w$. Suppose v_1 has no orbital intersecting A , in this case $w|_A = g|_A$, so that w would be an element of A_f which has a product decomposition using elements of Γ_1 with fewer uses of elements distinct from g_{11} and g_{11}^{-1} , which is impossible, by the construction of g . So v_1 has an orbital B_1 so that $B_1 \cap A \neq \emptyset$. Now, by the definition of the elements v_i , we know that the depth of B_1 is at least two in H_1 , so that $B_1 \subset A_2$. Define $C_1 = A$, and for each integer i where $2 \leq i \leq t$, inductively define $C_i = C_{i-1} v_{i-1}$. Suppose that for some $i \in \{1, 2, \dots, t\}$, $C_i \cap \text{Supp}(v_i) = \emptyset$. In this case, the product $v_1 v_2 \cdots v_{i-1} v_{i+1} \cdots v_t$, which is the product for g without the v_i term, will be a shorter product which equals f on A , which violates our construction. In particular, each C_i has non-trivial intersection with an orbital B_i of v_i . Now B_1 is an orbital of v_1 that has non-trivial intersection with $A = C_1$, and so $B_1 \subset A_2$, but now this implies that $C_2 \subset A_2$, and so $B_2 \subset A_2$. This same argument repeated $t - 2$ more times shows that for each $i \in \{1, 2, \dots, t\}$, we have $C_i \subset A_2$, and so $B_i \subset A_2$. In particular, each $v_i \in \Lambda_2$, or $v_i^{-1} \in \Lambda_2$. But now we have shown that g is a product of elements of H_2 , so our claim is proven.

Now define g_{21} to be a controller for H_2 on A_2 , and further define X_2 to be the set of towers for H_2 with largest element (A_2, g_{21}) . From the claim just proven (that H_2 admits towers of arbitrary height with largest elements of the form (A_2, h) for various $h \in H_2$), we see that X_2 has towers of arbitrary height.

Now define the following set:

$$\Gamma_2 = \{g \in H_2 \mid (g_{21}^k g) = \lambda, \lambda \in \Lambda_2, A_2 \not\subset \text{Supp}(g), k \in \mathbf{Z}\} \cup \{g_{21}\}$$

This is essentially looking at the interior parts of the generators of H_2 under the orbital A_2 of H_2 using the controller g_{21} .

It is immediate from the definition of controller that H_2 is generated by the finite set Γ_2 . Suppose that the order of Γ_2 is m_2 , and extend the indexing of g_{21} to the other elements of Γ_2 , so that $\Gamma_2 = \{g_{2i} \mid 1 \leq i \leq m_2, i \in \mathbf{N}\}$, where g_{21} is the only element of Γ_2 which realizes the orbital A_2 .

We now have that Γ_2 is a finite set of generators for H_2 which satisfies conditions (2) and (3).

Now by construction, H_2 , m_2 , Γ_2 , A_2 , and X_2 satisfy the six constraints.

But now, by temporarily relabeling all of the items mentioned in the six conditions for the series of groups $(H_i)_{i=1}^\infty$, with index one (the ones that all started with index two), we can repeat the argument that generated H_2 et al from H_1 , to create a new subgroup H_2 , and so forth. Remembering our original labelling, we see our new group and objects are H_3 , et al, which again satisfy conditions (1–4), (6). But now that A_3 satisfies condition (6), and A_2 satisfies condition (6) vis-a-vis A_1 , we see that X_3 satisfies condition (5). We can induct on this argument to now build the sequence of groups $(H_i)_{i=1}^\infty$ as promised, that satisfy all of the six conditions.

Let $T = \{(A_i, g_{i1}) | i \in \mathbf{N}\}$. By construction, $g_{i1} \in H_i \leq H$ has orbital A_i and $\bar{A}_{i+1} \subset A_i$ for all $i \in \mathbf{N}$. In particular, T is an infinite tower for H . $\diamond$

3.4 W and other groups

In this subsection, we will discuss how W relates to the various groups that we have found in the non-solvable subgroups of $PL_o(I)$. This will result in proofs of the Lemmas 3.6 and 3.5.

We first pick representations of the groups $(\mathbf{Z}l)^\infty$, $(\mathbf{Z})^\infty$, and $(\mathbf{Z}l)^\infty$ in $PL_o(I)$. Our presentations of these groups in Thompson's group F will be more explicit than that given in Brin's [2], and his proof that the resulting groups really are isomorphic with the wreath products named carries through with no difficulties, although in our concrete situation we can also use the outline of the proof in the introduction for at least the group $(\mathbf{Z})^\infty$.


Namely, define $\alpha \in PL_o(I)$ to be the element so that given any $x \in I$, we have

$$x\alpha = \begin{cases} \frac{1}{4}x & 0 \leq x < \frac{1}{4}, \\ x - \frac{3}{16} & \frac{1}{4} \leq x < \frac{7}{16}, \\ 4x - \frac{3}{2} & \frac{7}{16} \leq x < \frac{9}{16}, \\ x + \frac{3}{16} & \frac{9}{16} \leq x < \frac{3}{4}, \\ \frac{1}{4}x + \frac{3}{4} & \frac{3}{4} \leq x \leq 1, \end{cases}$$

and define $\beta_0 \in PL_o(I)$ to be the element so that given any $x \in I$, we have

$$x\beta_0 = \begin{cases} x & 0 \leq x < \frac{7}{16}, \\ 2x - \frac{7}{16} & \frac{7}{16} \leq x < \frac{15}{32}, \\ x + \frac{1}{32} & \frac{15}{32} \leq x < \frac{1}{2}, \\ \frac{1}{2}x + \frac{9}{32} & \frac{1}{2} \leq x < \frac{9}{16}, \\ x & \frac{9}{16} \leq x \leq 1. \end{cases}$$

The graphs of these elements (superimposed) are given below.



Now define $\beta_k = \beta_0^{\alpha^k}$ for each integer k . In particular, β_1 is given by the rule

$$x\beta_1 = x\alpha^{-1}\beta_0\alpha = \begin{cases} x & 0 \leq x < \frac{1}{4}, \\ 2x - \frac{1}{4} & \frac{1}{4} \leq x < \frac{3}{8}, \\ x + \frac{1}{8} & \frac{3}{8} \leq x < \frac{1}{2}, \\ \frac{1}{2}x + \frac{3}{8} & \frac{1}{2} \leq x < \frac{3}{4}, \\ x & \frac{3}{4} \leq x \leq 1. \end{cases}$$

Observe that the support of β_0 is $(\frac{7}{16}, \frac{9}{16})$, and since $\frac{7}{16}\beta_1 = \frac{9}{16}$, the support of β_0 is contained in a single fundamental domain of β_1 . In particular, given any $i \in \mathbf{Z}$, we see that the support of β_{i-1} is contained in a single fundamental domain of the support of β_i , since these two elements are conjugates of β_0 and β_1 . Now following Brin in [2], the group $(\imath\mathbf{Z})^\infty$ is the group generated by the full collection of the β_i , while $(\imath\mathbf{Z})^\infty$ is the group generated by the β_i with i a negative index, while $(\mathbf{Z}\imath)^\infty$ is the group generated by the β_i where we allow only the non-negative indices. The point is that whenever $i < j$ are indices, the support of β_i is fully contained inside a single fundamental domain of β_j . The reader may have realized that the group generated by the signatures of the tower found in Lemma 3.11 is isomorphic to $(\imath\mathbf{Z})^\infty$. Note that all of the β_i are found in Thompson's group F .

We will call β_{-1} the top generator of $(\imath\mathbf{Z})^\infty$, and β_0 the bottom generator of $(\mathbf{Z}\imath)^\infty$.

Lemma 3.19 *W embeds in both $(\imath\mathbf{Z})^\infty$ and $(\mathbf{Z}\imath)^\infty$.*

pf: We first embed W in $(\wr \mathbf{Z})^\infty$. For each $i \in \mathbf{N}$, define

$$\Gamma_i = \left\{ \gamma_j^i \mid \gamma_j^i = \beta_j^{\beta_{i+1}}, 1 \leq j \leq i, j \in \mathbf{N} \right\}$$

Note that each collection Γ_i generates a group isomorphic to W_i , by the argument given in the introduction after the discussion of W_2 , or also by the details of Brin in [2]. Further, the supports of the generators in Γ_i are all disjoint from the supports of the generators in Γ_j whenever $i, j \in \mathbf{N}$ with $i \neq j$, so that the elements of Γ_i found in $(\mathbf{Z}\wr)^\infty$ above commute with the elements of Γ_j in this case. Hence, the set

$$\Gamma = \cup_{i \in \mathbf{N}} \Gamma_i$$

generates a group

$$\langle \Gamma \rangle \cong \bigoplus_{i \in \mathbf{N}} W_i \cong W.$$

(Note: We will use this realization of W throughout the rest of this subsection when we refer to our realization of W in $PL_o(I)$. When we refer to “the first n summands of W ” we will mean the n subgroups (each isomorphic to a W_i) of W corresponding to the individual groups generated by the sets Γ_i for $1 \leq i \leq n, i \in \mathbf{N}$.)

We now embed W in $(\wr \mathbf{Z})^\infty$ in a similar fashion, finding copies of each W_i in $(\wr \mathbf{Z})^\infty$, all of which occur with mutually disjoint supports in I , the union of their generators will then generate a group isomorphic to W . Let $i \in \mathbf{N}$, and define

$$\Upsilon_i = \left\{ \theta_j^i \mid \theta_j^i = \beta_{-i+j-2}^{\beta_{-1}^{i-1}}, 1 \leq j \leq i, j \in \mathbf{N} \right\}.$$

so that Υ_i is the collection of the i generators beneath β_{-1} of the i 'th conjugate of the generators of $(\wr \mathbf{Z})^\infty$ by β_{-1} . Each collection Υ_i therefore generates a group isomorphic to W_i , while if $i \neq j$, any generator in Υ_i has disjoint support from the generators of Υ_j , so that the union

$$\Upsilon = \cup_{i \in \mathbf{N}} \Upsilon_i$$

has the property that

$$\langle \Upsilon \rangle \cong \bigoplus_{i \in \mathbf{N}} W_i \cong W.$$

◇

Lemma 3.20 *Neither $(\wr \mathbf{Z})^\infty$, nor $(\mathbf{Z}\wr)^\infty$, nor $(\wr \mathbf{Z}\wr)^\infty$ embed in W .*

pf:

If we show that one of $(\wr \mathbf{Z})^\infty$ or $(\mathbf{Z}\wr)^\infty$ fails to embed in W , then we will have shown that $(\wr \mathbf{Z}\wr)^\infty$ fails to embed in W , so we will say nothing more about $(\wr \mathbf{Z}\wr)^\infty$.

From the details of the proof of Lemma 2.28, we see that the orbital of a signed orbital in a tall tower of a subgroup G of $PL_o(I)$ survives to be an orbital of an element in the derived group G' . If we let $G^{(n)}$ temporarily represent the n 'th derived group of $(\mathbf{Z}\wr)^\infty$, as realized above, then each of the orbitals of the generators of $(\mathbf{Z}\wr)^\infty$ will be orbitals of some element of $G^{(n)}$. None of these elements of $G^{(n)}$ with orbitals shared with a generator β_k

(with $k > 0$, $k \in \mathbf{N}$) will commute with the generator β_0 in $(\mathbf{Z}l)^\infty$. Now suppose we have found an embedded copy of $(\mathbf{Z}l)^\infty$ in our realization of W above. There is $N \in \mathbf{N}$ so that the orbitals of the image of the bottom generator of $(\mathbf{Z}l)^\infty$ in W is contained in $\bigoplus_{1 \leq i \leq N} W_i$, the first N summands used in the definition of W , since elements of $PL_o(I)$ have finitely many orbitals. Now the N 'th derived group of W is trivial over the domain of these summands, by the argument of Lemma 2.30, so that the image of the bottom generator of $(\mathbf{Z}l)^\infty$ in W must commute with all of the elements of the M 'th derived group of $(\mathbf{Z}l)^\infty$ in W . Therefore, $(\mathbf{Z}l)^\infty$ cannot embed in W .

To see that $(\mathbf{Z})^\infty$ does not embed in W . Let $G^{(n)}$ now represent the n 'th derived group of $(\mathbf{Z})^\infty$. Suppose we have an embedding of $(\mathbf{Z})^\infty$ in W , and let α_{-1} represent the image of the top generator β_{-1} of $(\mathbf{Z})^\infty$ in W . Note that for any positive integer m , β_{-1} does not commute with any of the non-trivial elements of the $G^{(m)}$ in $(\mathbf{Z})^\infty$ and that this collection is non-empty. This implies that α_{-1} must not commute with some elements of the image of $G^{(m)}$ in W . Observe that this image is contained in $W^{(m)}$. Now suppose n is the largest index where the summand W_n of W shares some support with α_{-1} . α_{-1} must commute with every element of $W^{(n)}$ in W , since the group $W^{(n)}$ has no support in the supports of the first n summands of W , by Lemma 2.30, but this contradicts our statement that the subgroup $G^{(n)}$ of $W^{(n)} \leq W$ has elements that fail to commute with β_{-1} for any $n > 0$. $\diamond$

3.5 W in arbitrary non-solvable subgroups of $PL_o(I)$

The lemma below is a restatement of Lemma 3.7 and completes the proof of Theorem 1.2.

Lemma 3.21 *If H is a non-solvable subgroup of $PL_o(I)$ then H contains a subgroup isomorphic to W .*

pf:

Suppose H is a non-solvable subgroup of $PL_o(I)$. By Lemma 3.6 we know that W embeds in $(\mathbf{Z})^\infty$ and $(\mathbf{Z}l)^\infty$, and therefore also into $(\mathbf{Z}l)^\infty$. Therefore, if H admits infinite towers then we already have the result, so let us assume that H does not admit infinite towers. In particular, H admits towers of arbitrary finite height, H is balanced, and H admits no transition chains of length two.

Since H does not admit infinite towers, the depth of any signed orbital of H is well defined and finite. Since H is not the trivial group, H has a non-empty collection of orbitals. The analysis now breaks into two cases.

Case 1:

Suppose H admits no orbital that supports towers of arbitrary height. In this case the depth of any orbital of H is well defined, every orbital of H has finite depth, and given any $n \in \mathbf{N}$, H has orbitals with depth greater than n .

Now, pick an element g_1^1 of H so that $\hat{T}_1 = \{(B_1^1, g_1^1)\}$ is a tower of height one for H . g_1^1 will be our generator for W_1 . g_1^1 has finitely many orbitals, and so there is a maximum depth j_1 of the orbitals of H that are not disjoint from the support of g_1^1 . We will now pick our remaining generators from the group $H^{(j_1)}$, the j_1 'st derived subgroup of H . We note that no element in $H^{(j_1)}$ can have support intersecting g_1^1 , since $H^{(j_1)}$ has trivial support over the orbitals of H of depth less than or equal to j_1 as a consequence of the details of the proof of

Lemma 2.30. We also observe that $H^{(j_1)}$ still admits towers of arbitrary height, and infinitely many orbitals, of arbitrary finite depth. We now find a tower $\hat{T}_2 = \{(B_1^2, g_1^2), (B_2^2, g_2^2)\}$ for $H^{(j_1)} \leq H$ of height two. Now the signatures of $\hat{T}_2$ admit a finite total number of orbitals, and therefore the union of this collection of element orbitals is contained in the union of the collection of orbitals of H of depth less than some integer $j_2 > j_1$. We therefore will pick a tower $\hat{T}_3 = \{(B_1^3, g_1^3), (B_2^3, g_2^3), (B_3^3, g_3^3)\}$ for $H^{(j_2)}$ which has signatures whose supports must be disjoint from the supports of the signatures of the first two towers $\hat{T}_1$ and $\hat{T}_2$. We can continue in this fashion to inductively define towers $\hat{T}_k$ and integers j_{k-1} for each positive integer k so that the integers j_k are always getting larger, and so that the towers $\hat{T}_k$ always have height k and have signatures which are disjoint in support from the signatures of the previous towers. (This last follows since $\hat{T}_k$ has all of the orbitals of its signatures in orbitals of $H^{(j_{k-1})}$ whose supports are away from the orbitals of depth less than j_{k-1} of H , which orbitals contain all of the supports of the signatures of all of the towers with smaller index).

Let $k \in \mathbf{N}$. Let $\hat{G}_k$ represent the group generated by the signatures of $\hat{T}_k$. We can use the techniques of the proof of Lemma 3.17 to replace $\hat{T}_k$ with a new tower T_k supported by a subset of orbitals of H that support $\hat{T}_k$, so that the signatures of T_k generate a group G_k isomorphic to W_k . Do this for all $k \in \mathbf{N}$.

Now the union of all the signatures of all of the towers T_k forms a collection of generators of a group isomorphic to W .

Case 2:

Suppose now that H admits an orbital A that supports towers of arbitrary height.

If A is not an orbital of any element of H then A can be written as a union of an infinite collection of nested element orbitals of H , so that H would then admit an infinite tower, therefore there is an element d of H so that (A, d) is a signed orbital of depth one for H .

We will now restrict our attention to a special subgroup H_d of H which is directed by the element d , in a sense that will be made clear. Given any element $h \in H$, let k_h and j_h represent the smallest positive integers so that h^{k_h} and d^{j_h} satisfy the mutual efficiency condition. Let

$$\Gamma_d = \{[h^{k_h}, d^{j_h}] \mid h \in H\} \cup \{d\}.$$

The elements of Γ_d have all of their orbitals properly contained inside the orbitals of d , and since the orbital A of H admits towers of arbitrary height, and any element orbital B which is properly contained inside A will be realized as an orbital of some element g of Γ_d (note that it does not matter that we passed to high powers to guarantee the mutual efficiency condition), we see that the group $H_d = \langle \Gamma_d \rangle$ admits towers of arbitrary height. We now observe that given any finite set X of elements of H_d that do not support any signed orbitals of depth one for H_d , and a finite tower T for H_d which also contains no signed orbital of depth one, we can find a minimal power k_X of d so that so that the tower $T^{d^{k_X}}$ for H_d induced from T via conjugation of the signatures of T by d^{k_X} will have all of its signatures with disjoint support from the signatures of X , since we can conjugate the tower to be arbitrarily near to an end of an orbital of d . Therefore, for each positive integer n , let $\tilde{T}_n$ be a tower for H_d of height n . Now inductively define towers $\hat{T}_n$ which are towers induced from the $\tilde{T}_n$ by conjugation by powers of d so that given any positive integer k , the tower $\hat{T}_k$ has signatures whose supports are all disjoint from the signatures of the towers $\hat{T}_j$ whenever $j < k$ is a positive integer.

Now follow the procedure at the end of the previous case to improve the towers $\hat{T}_n$ to new towers T_n so that for each positive integer n , the signatures of the tower T_n generate a group isomorphic with W_n , while preserving the conditions that the signatures of distinct towers T_k and T_j have disjoint supports from each other, so that the union of all of the signatures of all of the towers T_k is a set of generators of a group isomorphic with W in H .

◇

References

- [1] Matthew G. Brin, *The ubiquity of Thompson's group F in groups of piecewise linear homeomorphisms of the unit interval*, J. London Math. Soc. (2) **60** (1999), no. 2, 449–460.
- [2] ———, *Elementary amenable subgroups of R . Thompson's group F* , Intl. J. Alg. and Comp. (to appear).
- [3] Matthew G. Brin and Craig C. Squier, *Presentations, conjugacy, roots, and centralizers in groups of piecewise linear homeomorphisms of the real line*, Comm. Algebra **29** (2001), no. 10, 4557–4596.
- [4] J. W. Cannon, W. J. Floyd, and W. R. Parry, *Introductory notes on Richard Thompson's groups*, Enseign. Math. (2) **42** (1996), no. 3–4, 215–256.
- [5] Andrés Navas, *Groupes résolubles de difféomorphismes de l'intervalle, du cercle et de la droite*, Bull. Braz. Math. Soc. (N.S.) **35** (2004), no. 1, 13–50.
- [6] ———, *Quelques groupes moyennables de difféomorphismes de l'intervalle*, Bol. Soc. Mat. Mexicana (3) **10** (2004), no. 2, 219–244 (2005).
- [7] Peter M. Neumann, *On the structure of standard wreath products of groups*, Math. Z. **84** (1964), 343–373.
- [8] Derek J. S. Robinson, *A course in the theory of groups*, second ed., Graduate Texts in Mathematics, vol. 80, Springer-Verlag, New York, 1996.